

# Carbohydrates in water-stable aggregates and particle size fractions of forested and cultivated soils in two contrasting tropical ecosystems

R. SPACCINI<sup>1</sup>, A. ZENA<sup>1</sup>, C.A. IGWE<sup>2</sup>, J.S.C. MBAGWU<sup>2</sup> & A. PICCOLO<sup>1\*</sup>

<sup>1</sup>Dipartimento di Scienze Chimico-Agrarie, Università di Napoli 'Federico II', Via Università 100, 80055 Portici, Italy; <sup>2</sup>Department of Soil Science, University of Nigeria, Nsukka, Nigeria (\*Author for correspondence, e-mail: alpiccol@unina.it; fax: 0039-081-7755130)

Key words: agroecosystems, carbohydrates, cultivation, forest soils, humid tropical soils, organic carbon, water-stable aggregates

Abstract. Information on changes in storage and loss of soil organic carbon (SOC) when tropical forests are converted to cropland is needed for evaluating soil structural degradation and for selecting appropriate sustainable soil management practices. We evaluated changes in SOC storage of organic carbon and acid-hydrolyzable carbohydrates content of aggregated classes and particle size fractions of adjacent forested and cultivated soils in eight agroecosystems from Ethiopian highlands and Nigerian lowlands. In all agroecosystems, SOC content was two to four times higher in the forested than the cultivated soils. Higher SOC content was found in Ethiopian  $(20.2-47.3 \text{ g.kg}^{-1})$  than Nigerian  $(12.0-24.0 \text{ g.kg}^{-1})$  forested soils. The magnitude of reduction in SOC and total carbohydrates with cultivation was soil-specific, being generally higher in the sandy than the clayey soils. The smaller aggregate classes (< 1.00 mm) and the sand-sized particles (2000-63  $\mu$ m) of the forested soils were preferentially enriched in carbohydrates relative to larger aggregates (4.75-1.00 mm). Carbohydrates were more concentrated in the clay-size fraction of the forested than in that of the cultivated soils. Cultivation reduced aggregate stability, increased the proportions of the smaller size aggregates and their associated carbohydrates relative to the forested soils. The susceptibility of the cultivated soils to loss in structural stability reflected this initial aggregation which was greater in the more stable clayey than the fragile sandy soils. The aggregate stability of either the forested or the cultivated soil could not be accounted for by the levels of OC or total carbohydrates in the soil.

# Introduction

Conversion of forests to arable lands is usually accompanied by a decline in soil organic carbon (SOC) content. The rate of such decline is influenced by intensity of cultivation, type of soil, texture and dominant mineralogy

(Mbagwu & Piccolo 1998) as well as by the ambient climate which is generally lower in cool temperate than in hot tropical environments (Arrouays & Pelissier 1994). Identification of the magnitude of such managementinduced SOC losses and of the most sensitive soil organic matter (SOM) pools affected is needed to select mitigating soil management practices (Adamu et al. 1997). Such identification is important for tropical regions where rates of deforestation are alarming (Lal 1986).

Soil carbohydrates which constitute from 5 to 25% of SOM (Stevenson 1994) are the main component of the labile pool of organic matter (OM) most affected by land use (Cambardella & Elliot 1992). Even though their long-term role in soil physical improvement is still controversial (Piccolo 1996; Degens & Sparling 1996), they are an important source of energy for soil microbial activity (Insam 1996). Extracellular gums and mucilages produced during microbial oxidation of OM are believed to temporarily stabilize soil aggregates (Tisdall & Oades 1982) and thereby protect them against deformation from heavy raindrop impacts which are common at the beginning of rains, especially in the tropics. Because of their labile nature, the effect of soil management practices on the concentrations of carbohydrates are greater than those on the more stable, recalcitrant humified fractions (Piccolo 1996).

Several studies have produced inconsistent results on the effects of land use on both total SOC and concentrations of carbohydrate fractions (Roberson et al. 1991; Dalal & Henry 1988; Arshad et al. 1990; Benzing-Purdie 1980). For example, Roberson et al. (1991) observed that after 2 years of conventional tillage in an orchard in California, soil carbohydrates were of the same level as those in the untilled plots. Arshad et al. (1990) and Hu et al. (1995) reported however, that long-term tillage reduced soil carbohydrates. Such differences in results may be related to texture, soil type, original soil carbon level and quality, and the distribution of natural aggregates in soils (Mbagwu & Piccolo 1998). In hot, humid southern Nigeria, Mbagwu and Piccolo (1998) observed that carbohydrates and SOC decreased with decreasing wet-aggregate diameter, whereas in the cool plateau region of central Nigeria, Adamu et al. (1997) using the same types of soils, reported that smaller dry aggregates were preferentially enriched in OC relative to the macroaggregates. This indicates possible confounding influences of environment and the type of aggregates studied (i.e., whether wet- or dry-sieved). In Canada, Smith et al. (1997) computed a total loss of 1.93 t.ha<sup>-1</sup> of SOC (equivalent to 7.08 t of CO<sub>2</sub>) from agricultural lands, with higher rates in coarse- than fine-textured soils. Chesire et al. (1990), in a long-term incubation study with <sup>14</sup>C-labelled barley, found that carbohydrate contents in size fractions were in the order: sand > silt > clay, which is similar to the results obtained by Angers and Mehuys (1990).

With few exceptions (Beare et al. 1994; Elliot 1986), most of these studies have concentrated on whole soil fractions (< 2.0 mm) and have focused on few soils within similar ecosystems. While this approach gives a generalized overview of the total carbohydrate content in soils, a proper understanding and modeling of SOM dynamics requires an evaluation of the locations of these labile OM pools within aggregates and particle size fractions. Their precise location gives an indication of their potential accessibility for microbial degradation (Cheshire & Mundie 1981; Angers & N'Dayegamiye 1991). To obtain more general results and to facilitate extrapolation of results, more extensive regional data are needed on the effects of land use, texture, elevation, climate and soil types on storage and loss of OC when forests are converted to permanent arable farming (Davidson & Ackerman 1993; Arrouays & Pelissier 1994).

This paper presents the results of our study on the distribution of acidhydrolyzable carbohydrates in whole soils (< 2.0 mm diameter), water-stable aggregates and particle size fractions of forested and cultivated soils in different agroecosystems in sub-saharan Africa. The objectives were (1) to evaluate changes in carbohydrate concentrations in whole soils, aggregates and particle size fractions of soils before and after deforestation and subsequent arable farming and (2) to relate such changes to aggregate stability of the soils.

# Materials and methods

# Description of sites, experimental design and soil sampling

This study was conducted with soils from low-input agricultural systems in Ethiopia and southern Nigeria. This low-input agricultural system is such that forested sites are cleared for arable cropping without further inputs of organic or inorganic fertilizers. Also cultivation consists essentially of hand hoeing and planting without mechanical tillage in all sites but Holeta, Jimma and Umudike where mechanical cultivation (disking, ploughing and harrowing) was used. Relevant information on the sites is given in Table 1.

In each location the experimental set up was a Randomized Complete Block Design with each treatment replicated three times. The treatments were forested and cultivated sites. In Ethiopia, each plot size measured 6 m × 4 m, whereas in Nigeria each plot size was 6 m × 5 m. From September to November 1995, three bulked topsoil (0–20 cm) samples, collected from each replicate of the forested and cultivated sites in these locations, were airdried, sieved through 4.75 mm mesh and shipped to Italy for determination of water-stable aggregates and carbohydrates. The < 2.00 mm (fine earth) Table 1. Locations soils and land use histories of the sites studied.

Locations	Land use histories			
on of the locations of	ETHIOPIA			
Awassa:				

06° 58'N, 38° 25'E; Alt. 1700 m.a.s.l. Soil: Andisol; Annual rainfall: 1100 mm. Drainage class: Rapid.

Parent Material: Volcanic Ash. Vegetation: Highland Savannah

### Ginchi:

09° 01'N, 38°20'E; Alt. 2300 m.a.s.l. Soil: Vertisol; Annual rainfall: 900 mm. Drainage class: Moderate Parent Material: Weathered Basalt Vegetation: Secondary Forest

### Holeta:

09° 03'N, 38°30'E; Alt. 2400 m.a.s.l. Soil: Vertisol; Annual rainfall: 1054 mm Drainage class: Rapid Parent Material: Olivine Basalt Vegetation: Secondary Forest

### Jimma:

07° 41'N, 36° 50'E; Alt. 1800 m.a.s.l. Soil: Alfisol; Annual rainfall: 1400 mm Drainage class: Rapid Parent Material: Basalt Vegetation: Rainforest

### Sirinka:

11° 62'N, 39° 70'E; Alt. 1900 m.a.s.l. Soil: Entisol; Annual rainfall: 700 mm Drainage class: Moderate Parent Material: Alluvial Deposits Vegetation: Bush regrowths Located at the centre of the Rift Valley in the southwest part of Ethiopia; climax vegetation which represents the forested soil is highland savannah whereas the cultivated soil has been cropped for 4 years (1992–1995) and grown annually to maize.

Located at the central highlands; forest site is a secondary forest not cultivated for < 35 years and the cultivated soil has been planted to wheat and barley for 4 years (1992–1995)

Also located in the central highlands with vegetation similar to Ghinchi. Forested soil is a secondary regrowth not cultivated for about 20 years; cultivated soil has been grown to wheat and barley for 4 years (1992–1995).

Located in the western part with rainforest vegetation as the climax vegetation. The cultivated soil has been cropped to maize for 5 years (1988–1992) without returning maize residues to the soil.

Located in the highly degraded Tigrai region of northern Ethiopia where the climax vegetation is bush regrowths no cultivated for about 15 years. The cultivated soil has been cropped continuously to sorghum for 5 years (1988– 1998)

### Table 1 Continued

Locations	Land use histories

### NIGERIA

### Abakiliki.

06° 30'N, 08° 15'E, Alt. 200 m.a.s.l. Soil: Gravelly Inceptisol; Annual rainfall: 1560 mm Drainage class: Slow Parent Material: Shale Deposits Vegetation: Derived Savannah

#### Nsukka.

06° 51'N, 07° 24'E, Alt. 400 m.a.s.l. Soil: Ultisol; Annual rainfall: 1600 mm Drainage class: Very rapid Parent Material: False-bedded Sandstones Vegetation: Derived Savannah

# Forested soil is revegetated with Gmelina trees (established in 1960). Cultivated soil site is continuously cultivated for 5 years (1991-1995) with maize-vegetables-upland rice/soybeanmaize/cowpea rotation.

Forested soil represents the climax vegetation not affected by annual bush burning. Cultivated soil has been cultivated annually to cassava for 5 years (1991-1995).

### Umudike:

05° 29'N, 07° 33'E, Alt. 120 m.a.s.l. Drainage class: Very rapid Parent Material: Sedimentary Rocks Vegetation: Humid Rainforest

Forested soil is climax vegetation (> 100 years Soil: Ultisol, Annual rainfall: 2100 mm old) and the cultivated soil has been continuously cropped to cassava for 15 years.

fractions were used to measure particle size distributions (after complete dispersion with sodium hexametaphosphate) by the pipette method of Gee and Bauder (1986). pH was measured in 1:2.5 soil-water ratio, OC by the wet oxidation procedure of Nelson and Sommers (1982), and NH<sub>4</sub>-acetate exchangeable bases and CEC following procedures outlined in Anderson and Ingram (1993). Total carbohydrate content was measured in this whole soil fraction by the phenol-sulphuric acid method described below (Piccolo et al. 1996). All determinations were made on each of the replicated samples. The pertinent soil properties shown in Table 2 indicate significant redistribution between silt and clay for Holeta, Jimma, and Umudike due apparently to the use of mechanical cultivation (which involves soil inversion) in the cultivated sites and wind erosion effects.

Location/	Sand	Slit	Clay	pH	$TEB^1$	CEC <sup>2</sup>
land use	(g/kg)	(g/kg)	(g/kg)	(H <sub>2</sub> O)	(Cmol/kg)	(Cmol/kg)
er, ar sus bes	landint?	EI	THIOPIAN	SOILS	and a standard	Al (asymbolic
Awassa						
F	425	300	275	5.28	16.29	30.4
С	425	300	275	5.45	11.36	22.6
Ginchi						
F	25	263	712	5.74	29.41	54.4
С	25	263	712	5.98	29.11	54.0
Holeta						
F	57	238	705	5.58	10.55	31.8
С	0	163	837	4.88	4.12	29.6
Jimma						
F	100	250	650	6.15	16.07	30.2
С	25	143	823	4.94	7.74	26.2
Sirinka						
F	225	250	525	6.02	34.61	56.0
С	275	212	513	6.75	25.40	60.0
			ICEDIAN	COT C		
		N	IGERIAN	SOILS		
Abakiliki						
F	540	200	260	5.59	16.40	34.5
С	580	180	240	5.54	7.23	23.0
Nsukka						
F	740	40	220	3.53	0.99	15.0
С	700	20	280	3.87	4.27	14.5
Umudike						
F	660	60	280	4.21	6.93	24.0
С	820	20	160	4.53	1.84	11.0

Table 2. Some properties of the 0-20 cm of the forested (F) and cultivated (C) soils.

<sup>1</sup>TEB is total exchangeable bases.

<sup>2</sup>CEC is Cation exchange Capacity.

# Separation of water-stable aggregates and determination of aggregate stability

The method of Kemper and Rosenau (1986) was used to separate water-stable aggregates. Twenty grams of the < 4.75 mm air-dried soil were put in the topmost of a nest of three sieves of 1.00, 0.50, and 0.25 mm mesh size and pre-soaked in distilled water for 30 min. Thereafter the nest of sieves and its contents were oscillated vertically in water 20 times using a 4 cm amplitude at the rate of one oscillation per s. Care was taken to ensure that the soil particles on the topmost sieve were always below the water surface during each oscillation. After wet-sieving, the resistant soil materials on each sieve and the unstable (< 0.25 mm) aggregates were quantitatively transferred into beakers, dried in the oven at 50 °C for 48 h, weighed and stored for analysis of carbohydrates. The percentage ratio of the aggregates in each sieve represents the water-stable aggregates (WSA) of size classes: 4.75-1.00 mm, 1.00-0.50 mm, 0.50–0.25 mm and < 0.25 mm. Due to our interest in also studying the carbohydrate distribution in the particle size fractions we did not correct the WSA for the sand particles that may have co-settled with the aggregates on the sieves. Aggregate stability was measured as the mean-weight diameter (MWD) of water-stable aggregates as by equation 1:

$$MWD = \sum_{i=1}^{n} X_i W_i \tag{1}$$

where,  $X_i$  is the mean diameter of the i<sup>th</sup> sieve size and  $W_i$  is the proportion of the total aggregates in the I<sup>th</sup> fraction. Higher MWD values indicate higher proportions of macroaggregates in the sample and therefore, higher stability.

### Separation of particle size fractions

The coarse sand (2000–200  $\mu$ m), fine sand (200–63  $\mu$ m), silt (63–2  $\mu$ m) and clay (< 2  $\mu$ m)-sized fractions were separated from the < 2.00 mm particles without prior chemical pretreatment. Low sonication energy was used to disperse the soils so as to simulate natural disruptive forces such as from raindrop impact and avoid organic matter redistribution among the separates. The procedure (carried out in triplicate per sample) is the one described by Stemmer et al. (1997) and consisted of suspending 40 g of the < 2.0 mm, air-dried sample in 100 mL of deionized water in a beaker and dispersing with an ultrasonic probe (Ultrasonic Liquid Processor, XL-Series Sonicator) placed at about 1 cm depth from the soil-water suspension. The suspension was sonicated for 2 min yielding a total energy of 240 J (or 2.40 J mL<sup>-1</sup>). The dispersed soil was then wet-sieved through a series of sieves to separate

the coarse sand and fine sand. The silt fraction was separated through centrifugation at 90 g for 6 min. To maximize the silt yield, centrifugation was repeated two more times on the residue. The clay fraction was separated from the supernatant after centrifugation at 3100 g for 30 min. These fractions were air-dried, weighed and stored for determination of total carbohydrates. In both determinations of WSA and particle size fractions variation within triplicates of reference samples was less than 5%.

# Determination of total carbohydrates

For more detailed study of the distribution of carbohydrates within these samples, carbohydrate content was measured in duplicate per sample on the water-stable aggregates and particle-size fractions (separated as described above) by first hydrolyzing 1 g of soil by shaking for 16 h with 10 mL of a 0.25 M H<sub>2</sub>SO<sub>4</sub> solution in a rotary shaker. Interfering ions in the hydrolyzate were reduced by elution through anion and cation exchange resins (Martens & Frankenberger 1993). The monosaccharide content in the hydrolyzates was measured colorimetrically as glucose equivalents using the phenol-sulphuric acid method (Piccolo et al. 1996). All measurements were expressed as glucose concentrations in g.kg<sup>-1</sup> of the water-stable aggregates or the particle-size fractions.

### Data analysis

Simple correlations between MWD and SOC or total carbohydrates were used to evaluate their possible contribution to soil structural stability. For each site an analysis of variance (ANOVA) for Randomized Complete Block Design (RCBD) was used to compare the effects of land use on the measured properties. Where the F-values were significant at P = 0.05 level, the least significant differences (LSD) were calculated and used to compare the mean values of the forested and cultivated treatments. For ease of comparing the effects of land use on aggregate stability (AS) across all locations, the change in MWD between forest and cultivated sites was normalized as follows:

$$AS_{\nu} = \{1 - (MWD_c/MWD_f)\} \times 100$$
<sup>(2)</sup>

where,  $AS_{\nu}$  is the normalized MWD and subscripts *c* and *f* refer to cultivated and forested sites, respectively. In this way, comparison of the changes in structural stability and in SOC and carbohydrates across all locations becomes valid (Mbagwu et al. 1991).

### Results

# Effects of cultivation on water-stable aggregates and aggregate stability

Table 3 shows that irrespective of site, the proportion of the largest aggregate class (4.75–1.00 mm) decreased with cultivation. In Ethiopia cultivation reduced the proportion of water-stable aggregates between 51 and 97% relative to the five forested soils. In Nigeria, cultivation reduced the proportion of the water-stable aggregates between 6 and 61% relative to the three forested soils, showing a generally lower impact of cultivation on the sandy than clayey soils. Also with the exception of the gravelly soil from Abakiliki, soils with initial high percent of water-stable macroaggregates tended to be more susceptible to reduction in this aggregate-size range with cultivation.

At each site in Ethiopia, cultivation increased substantially the proportion of < 1.00 mm aggregate class. The average increases ranged from 45% to 70% for the forested sites and from 85% to 96% for the cultivated sites. The Vertisols were the most susceptible to the cultivation-related increase in small (< 1.00 mm) aggregates. In Nigeria the average forested and cultivated values were 67% and 75%, respectively. Hence deforestation and intensive arable farming on these soils reduced water-stable aggregates to smaller diameters. This was also reflected in the aggregate stability index used (MWD) which showed consistently higher values in the forested than in cultivated sites (Table 4). In Ethiopia the average reductions in MWD were 76% on the Vertisols, 63% on the Alfisol, 60% on the Entisol and 37% on the Andisol (Table 5). Reductions in MWD of the cultivated soils were less in sandier Nigerian than Ethiopian soils, and ranged from 7 to 22%.

### Effects of cultivation on organic carbon and total carbohydrate contents

In Ethiopian forested soils (Table 4), OC ranged from 27.2 g kg<sup>-1</sup> (in Sirinka) to 47.3 g kg<sup>-1</sup> (in Jimma), whereas OC content in Nigerian forested soils varied between 12.0 g kg<sup>-1</sup> (in Nsukka) and 24.0 g kg<sup>-1</sup> (in Abakiliki). The OC in cultivated sites was 50 to 75% lower than in forested sites. Similar results were obtained for total carbohydrates. Relative to the forested soils the largest reductions in OC occurred in the cultivated sandy Sirinka (72%) and Umudike (76%) soils (Table 5). The surface soil at these locations also had the largest reductions in carbohydrate content (65% and 75%, respectively).

Location/	Size fractions of WSA (mm)				
land use	4.75-1.00	1.00-0.50	0.50-0.25	< 0.25	
A SPACE AND A SPACE AND A	ETH	OPIAN SOIL	S	within	
Awassa					
F	30.3	19.2	19.8	30.7	
С	14.8	29.0	25.0	31.2	
LSD (0.05)	5.6	6.1	3.8	NS	
Ginchi					
F	47.4	10.7	7.0	34.9	
С	7.2	11.6	18.0	63.2	
LSD (0.05)	9.8	NS	5.5	11.4	
Holeta					
F	47.4	16.4	11.7	24.2	
С	1.2	7.0	31.8	60.0	
LSD (0.05)	10.3	4.5	8.2	11.1	
Jimma					
F	53.1	15.0	11.8	20.1	
С	13.0	21.5	29.0	36.5	
LSD (0.05)	8.9	3.3	8.8	9.3	
Sirinka					
F	55.3	18.2	7.7	18.8	
С	11.7	39.4	24.5	24.4	
LSD (0.05)	10.9	6.6	8.6	NS	
	NIG	ERIAN SOILS	nographing the		
Abakiliki					
F	63.0	13.6	7.0	16.4	
С	59.2	9.0	6.1	25.7	
LSD (0.05)	NS	2.3	NS	4.4	
Nsukka					
F	14.7	33.2	26.7	25.4	
С	5.8	36.4	37.9	19.9	
LSD (0.05)	3.5	NS	9.1	4.9	

Table 3. Distribution of water-stable aggregates (WSA, %) in forested (F) and cultivated (C) soils in Ethiopia and Nigeria.

Tabl	e 3.	Continued.

Location/	Size fractions of WSA (mm)					
land use	4.75-1.00	1.00-0.50	0.50-0.25	< 0.25		
Umudike	200-01	in the second projection	And Market States, Colors	CLASS V.		
F	19.8	37.8	19.1	23.3		
С	8.8	58.1	15.8	17.3		
LSD (0.05)	7.2	11.5	NS	4.8		

NS = Not significant.

The regression equations relating aggregate stability (AS) with OC and carbohydrates for all the Ethiopian and Nigerian forested (F) or cultivated (C) soils are:

AS(F) = 0.827 + 0.013(OC)	$(R^2 = 0.13)(N = 8)$	(3)
AS(C) = 0.958 - 0.023(OC)	$(R^2 = 0.09)(N = 8)$	(4)
AS(F) = 0.808 + 0.031(Carbohydrates)	$(R^2 = 0.23)(N = 8)$	(5)
AS(C) = 0.381 + 0.031(Carbohydrates)	$(R^2 = 0.09)(N = 8)$	(6)

None of the  $R^2$  values in eqs. 3–6 are significant, explaining only 9 to 13% of variation in AS with the level of OC present and between 9% and 23% of variation in AS with the level of carbohydrates in soils. Possible explanations for these low correlations are (i) that other SOC fractions are responsible for stabilizing aggregates in these soils (Monreal et al. 1995; Dinel et al. 1997; Piccolo & Mbagwu 1999), and (ii) that reduction in aggregates to smaller sizes due to tillage implements.

# Distribution of total carbohydrates in water-stable aggregates

In the forested sites, the smaller aggregate classes (< 1.00 mm) were preferentially enriched in total carbohydrates as compared with the large (4.75–1.00 mm) aggregates (Table 6). The carbohydrates stored in the small (< 1.00 mm) aggregates of the forested soils ranged from 7.2 g kg<sup>-1</sup> (Nsukka) to 16.6 g kg<sup>-1</sup> in Awassa whereas the concentrations of those stored in the macroaggregate classes (> 1.00 mm) were lower and varied between 2.0 g kg<sup>-1</sup> (Nsukka) and 5.3 g kg<sup>-1</sup> (Abakiliki) (Table 6). This indicates that the carbohydrates are better protected in the smaller than in the larger size aggregates of the forested soils. With cultivation the concentration of carbohydrates in these smaller class aggregates was reduced in absolute values but were still

Location/ land use	Aggregate Stability (MWD, mm)	OC (g/kg)	Total carbohydrates (g/kg)	Proportion of carbohydrates in < 0.25 mm, (%)	Carbohydrates in (< 1.0 mm)/ Carbohydrates in (> 1.0 mm)
	ing grat	Е	THIOPIAN SOI	LS	150%
Awassa					
F	0.928	40.3	21.0	33.7	3.73
С	0.582	16.6	14.1	22.5	2.39
LSD (0.05)	0.105	6.5	3.54	7.18	0.97
Ginchi					
F	1.324	32.8	13.40	19.8	4.09
С	0.362	20.0	9.16	43.8	2.52
LSD (0.05)	0.244	8.3	2.45	2.20	1.10
Holeta					
F	1.342	36.1	14.64	13.6	3.85
C	0.267	16.0	9.88	23.2	0.90
LSD (0.05)	0.112	10.3	3.68	3.82	2.05
Jimma					
F	1.491	47.3	12.40	19.1	3.16
С	0.545	20.2	9.28	22.6	3.03
LSD (0.05)	0.234	11.9	2.06	2.75	0.15
Sirinka					
F	1.443	27.2	12.44	44.3	3.08
С	0.582	7.7	4.36	47.9	0.33
LSD (0.05)	0.199	3.1	5.72	3.8	2.00
		huble a	UCEDIAN COU	Sharhodnas lord	Distribution, of th
41 1.1.1.1.			NIGERIAN SUI	una multi protia i	
Ε	1 070	24.0	16.22	50.1	2.00
F	1.8/8	10.4	10.32	50.1	2.09
	1./41	10.4	2.39	7.70	1.93
LSD (0.05)	142	1.2	2.87	7.70	0.17
Nsukka					
F	0.621	12.0	6.90	50.4	2.38
С	0.527	6.8	3.27	54.1	0.01
LSD (0.05)	0.065	2.1	2.65	4.6	1.75

*Table 4.* Differences in aggregate stability, organic carbon (OC) and total carbohydrates of forested (F) and cultivated (C) soils in Ethiopia and Nigeria.

Table 4. Continued.

Location/ land use	Aggregate Stability (MWD, mm)	OC (g/kg)	Total carbohydrates (g/kg)	Proportion of carbohydrates in < 0.25 mm, (%)	Carbohydrates in (< 1.0 mm)/ Carbohydrates in (> 1.0 mm)
Umudike		5.445	HORMAN SOL	18. 0.78	1.05
F	0.826	23.6	12.81	15.7	2.63
С	0.648	5.6	3.25	32.3	3.06
LSD (0.05)	0.105	3.0	6.58	11.38	0.30

*Table 5.* Normalized changes  $(\%)^1$  in soil organic carbon (OC), total carbohydrates and aggregate stability from forested to cultivated soils in Ethiopia and Nigeria.

Locations	OC	Total carbohydrates	Aggregate stability
testellar of lan	ETHIO	PIAN SOILS	tions Jectim
Awassa	58.8	33.0	37.3
Ginchi	39.0	31.2	72.7
Holeta	55.7	29.8	80.1
Jimma	57.3	25.2	63.4
Sirinka	71.7	65.0	59.7
	NIGER	IAN SOILS	
Abakiliki	56.7	24.1	7.3
Nsukka	43.3	52.6	15.1
Umudike	76.3	74.6	21.5

<sup>1</sup>Normalized value =  $[1 - (cultivated-value/forested-value)] \times 100$ . All values indicate percentage reduction relative to forests.

better protected in these aggregates than in the larger ones. The ratio of carbohydrates in the < 1.00 mm aggregates to that in the > 1.00 mm aggregates of the forested soils varied between 2.1 to 4.1, thereby indicating a two- to four-fold increase in carbohydrates stabilization.

In all aggregate sizes, cultivation reduced carbohydrate concentrations (Table 6). Hence the ratios of their distribution in the < 1.00 mm to the > 1.00 mm aggregates were generally lower in the cultivated than in the forested soils (Table 4). Also the proportion of carbohydrates in the < 0.25 mm aggregates of the forested soils varied between 13.6% (in Holeta)

Location/	Water Stable Aggregate Classes (mm)					
land use	4.75-1.00	1.00-0.50	0.50-0.25	< 0.25		
	ETH	IIOPIAN SOILS	5	Col 10 and		
Awassa						
F and	4.44	4.58	4.92	7.07		
С	4.18	4.40	2.35	3.17		
LSD (0.05)	0.27	0.09	1.27	1.59		
Ghinchi						
attai <mark>r</mark> dodaa hool	2.64	2.29	1.58	2.65		
C	2.60	2.81	3.60	4.01		
LSD (0.05)	0.07	0.37	1.48	0.96		
Holeta						
F	3.02	2.47	2.41	1.99		
С	5.20	3.34	3.82	2.29		
LSD (0.05)	1.45	0.56	1.00	0.21		
Jimma						
F	2.98	4.33	2.72	2.37		
С	2.30	2.46	2.43	2.10		
LSD (0.05)	0.48	1.32	0.42	0.20		
Sirinka						
F	3.05	1.87	1.99	5.51		
С	3.27	1.30	1.69	2.09		
LSD (0.05)	0.26	0.43	0.22	2.42		
	NIC	BERIAN SOILS	ic) of Stituler of			
Abakiliki						
F	5.28	10.00	9.16	8.17		
С	4.23	7.30	5.80	7.45		
LSD (0.05)	0.47	1.83	2.75	0.91		
Nsukka						
Find and	2.04	1.92	1.77	3.48		
C	3.23	2.02	1.75	1.77		
LSD (0.05)	0.84	0.17	0.04	1.21		

*Table 6.* Concentrations of carbohydrates (g/kg) in water-stable aggregate (WSA) classes of forested (F) and Cultivated (C) soils in Ethiopia and Nigeria.

Location/	Water Stable Aggregate Classes (mm)					
land use	4.75-1.00	1.00-0.50	0.50-0.25	< 0.25		
Umudike	(m) 20-0000	0 0mg 50-00	0-206 pm) (0	ion -		
F	3.53	1.91	1.86	2.01		
С	0.84	0.87	0.78	1.05		
LSD (0.05)	1.39	0.75	0.76	0.69		

Table 6. Continued.

and 50.4% (in Abakiliki) whereas in the cultivated soils it ranged from 22.5% (in Awassa) to 60.1% (in Abakiliki). This trend shows, therefore, that as forests are converted to arable lands, there is a preferential accumulation of carbohydrates in small aggregates < 1.00 mm diameter (Table 4).

# Distribution of carbohydrates in particle size fractions

Irrespective of land use, the highest carbohydrate concentrations occurred in the total sand-sized fraction (2000–63  $\mu$ m) (Table 7). Most of the forested, clay-sized fraction accumulated more carbohydrates than the silt-sized fraction whereas in most of the cultivated soils the silt-sized fraction contained more carbohydrates than the clay-sized fraction. This shows that in most sites, carbohydrates are protected more in clay-sized fractions under forest than under cultivation apparently because the clay-sized fractions have large surface areas and can adsorb carbohydrates which may be further protected in strong clay-OM complexes.

In three of the Ethiopian sites (Ginchi, Holeta and Sirinka), cultivation increased the concentrations of carbohydrates in the clay fraction while reducing them in the sand fraction as compared to forested soils (Table 7). No significant change was observed between the carbohydrate concentrations in the clay fractions of forested and cultivated Awassa soils. Cultivation decreased carbohydrate concentrations in all particle size fractions of Jimma soil. In Nigeria, cultivation also decreased carbohydrates in all size fractions at Abakiliki and only in the sand and clay fractions at Umudike (Table 7). At Nsukka, cultivation caused an increase in soil carbohydrate concentrations consistently in all size fractions.

Location/ land use	Coarse sand (2000–200 μm)	Fine sand (200–63 μm)	Total sand (2000–63 μm)	Silt (63–2 μm)	Clay (< 2 μm)
Awassa					
F	3.54	3.92	7.46	5.12	2.19
С	2.25	2.37	4.62	1.97	2.01
LSD (0.05)	0.81	0.85	1.48	1.56	NS
Ghinchi					
F	2.00	2.04	4.04	0.35	0.95
С	0.75	3.00	3.75	1.57	1.36
LSD (0.05)	0.63	0.71	NS	0.83	0.22
Holeta					
F	2.99	2.90	5.89	2.50	1.01
С	1.69	2.93	4.62	3.14	1.89
LDS (0.05)	0.42	NS	0.74	0.32	0.51
Jimma					
F	5.47	4.23	9.70	4.96	5.33
С	4.81	3.19	8.00	3.91	3.03
LSD (0.05)	0.54	0.62	1.32	0.85	1.09
Sirinka					
F	3.27	4.53	7.80	4.26	4.45
С	5.70	7.30	13.00	6.92	4.93
LSD (0.05)	1.30	1.56	3.46	1.77	NS
		NICEDIAN	0.000		
Abakiliki		NIGERIAN	SOILS		
F	3 56	4 46	8.02	3.42	4.00
C	3.74	2.94	6.68	1.59	3.13
LSD (0.05)	NS	0.65	1.16	0.92	0.60
Nsukka					
F	0.87	1.64	2.51	2.24	0.96
С	1.08	1.74	2.80	2.50	1.40
LSD (0.05)	0.08	NS	0.26	0.20	0.28

*Table 7.* Concentrations of carbohydrates (g/kg) in particle size fractions of forested (F) and cultivated (C) soils in Ethiopia and Nigeria.

Location/ land use	Coarse sand (2000–200 µm)	Fine sand (200–63 µm)	Total sand (2000–63 μm)	Silt (63–2 μm)	Clay (< 2 μm)
Umudike	We dist he will	wide Fastin	NUMBER OF STREET	ande kåb	Carrie Like
F	1.26	1.37	2.63	1.39	2.23
С	0.62	1.64	2.26	1.61	2.14
LSD (0.05)	0.39	0.21	0.29	0.18	NS
	T PATRO AND AND AND AND	The second second second			

Table 7. Continued.

NS = Not significant.

### Discussion

We concentrated on effects of deforestation and subsequent cultivation on soil organic matter storage, quality, and aggregate stability. In Africa, this low-input system represents an extensive worst-case scenario for studying SOM and aggregate degradation processes arising from long-term continuous cultivation. Our observation of reductions in the proportion of macroaggregates (> 0.25 mm) following cultivation was also reported in other ecosystems. For example, on a sandy clay loam from Georgia, USA, Beare et al. (1994) obtained higher proportion of macroaggregates under zero than conventional-tillage practices. Cambardella and Elliot (1993) and Gupta and Germida (1988) reported higher proportions of microaggregates (< 0.25 mm) when native grasslands were cultivated. The greater reduction in proportion of the largest aggregate size (4.75-1.00 mm) obtained in Ethiopia than Nigeria may be due to the generally higher initial aggregation of Ethiopia compared to Nigeria forested soils. In Ethiopia, the average proportion of < 1.00 mm aggregates in the cultivated soils varied in the order: Holeta (99%) > Ginchi (93%) > Sirinka (88%) > Jimma (87%) > Awassa (85%) and in Nigeria, Nsukka (94%) > Umudike (91%) > Abakiliki (41%). The latter shows that the Ethiopian Vertisols and the Nigerian Ultisols are more susceptible to reduction in the proportion of larger aggregates following cultivation than in other soils.

The decrease in total SOC and carbohydrates in forested soils with cultivation is attributed to disintegration of the larger aggregates and consequent exposure to microbial oxidation of the partially decomposed organic materials previously protected in the disintegrated aggregates (Dalal & Henry 1988; Cambardella & Elliot 1993). Roberson et al. (1991) reported earlier that tillage had no significant effect on the OC content of a loamy soil from California (USA) which is the reverse of our observation. A number of papers (Hu et al. 1995, and Salinas-Garcia 1997 in USA; Chan 1997 in Australia; Arshad et al. 1990; Baldock et al. 1987; Angers & Mehuys 1989, and Franzluebbers & Arshad 1997 in Canada and Mbagwu & Piccolo, 1998 in Nigeria) reported results similar to those of our study. Recently Parfitt et al. (1997) related the stability and turnover time of OC with mineralogy of Al and Fe oxides by suggesting that SOC was stabilized more in soils with higher amount of allophane and ferrihydrite (eg. Andisols) than in other soils. Monreal et al. (1997) and Monreal and Kodama (1997) while agreeing with the findings of Parfitt et al. (1997) for OC, noted that the carbohydrate fraction was not affected by such abiotic stabilization reactions with the oxides of Fe and Al. In our study the percentage loss of OC in the Andisol (Awassa) was the second highest in Ethiopia, which indicates no such preferential stabilization of OC in this Andisol.

The general trend of the smaller forest aggregates to be preferentially enriched in total carbohydrates was also reported by Webber (1965). A possible explanation could be that carbohydrates in smaller aggregates are more strongly adsorbed as polymers of various forms and dimensions to either clay or humic fractions which abound in the microaggregates (Stevenson 1994; Piccolo & Mbagwu 1990). In this way the carbohydrates are better protected from microbial degradation than those in the larger aggregates, hence their higher concentrations in the smaller aggregate fractions. The cultivation-induced reduction of carbohydrates in aggregates > 1.00 mm and their corresponding increase in aggregates < 1.00 mm of most of these soils supports the findings of Cambardella and Elliot (1993). Likewise, Dalal and Henry (1988) proposed that destruction of macroaggregates by cultivation exposes physically protected organic materials to rapid oxidation and loss. This hypothesis was later confirmed by Monreal et al. (1997) who showed that SOC in macroaggregates > 250  $\mu$ m was young and turned over in 14 years.

Irrespective of site and cultivation history, the total sand fraction had the highest concentration of carbohydrates. Loss of carbohydrates from sands following cultivation varied from 7% in clayey Ginchi to 38% in sandy Awassa. During an initial period of cultivation, Dalal and Mayer (1986) also reported rapid loss of sand-sized OM (which was equivalent to their particulate OM fraction).

The lack of a strong relationship between aggregate stability and OC or carbohydrates in this study suggests that other binding agents may be involved in stabilizing the aggregates in these soils. Carbohydrates have been previously associated with improvements in soil aggregate stability (Angers & Mehuys 1989; Haynes & Swift 1990; Angers et al. 1993; Haynes & Francis 1993). However, according to Insam (1996), since carbohydrates are easily degraded by microorganisms, they cannot participate in long-term stabiliz-

ation of soil aggregates. The short term effect of a polysaccharide on soil aggregate stability was recently confirmed by Piccolo and Mbagwu (1999). Dutarte et al. (1993) associated aggregate stability on intensively cultivated sandy soils from West Africa with the humified fractions of organic matter, notably humin and humic acids. Earlier studies (Piccolo & Mbagwu 1990; Mbagwu & Piccolo 1998) also reported a significantly higher positive correlation between microaggregate stability and humic acid content than either total OC or the carbohydrate contents. Carbohydrates have been shown to participate in the process of aggregate stabilization when acting in conjunction with the more humified soil organic matter pools (Angers & Mehuys 1989; Caron et al. 1992; Piccolo & Mbagwu 1999). Hence isolating carbohydrates alone and testing statistically their effect on aggregate stability indicates weak relationships as observed in this study.

In conclusion, these results show that continuous cultivation following deforestation reduced the proportion of the largest aggregate class (4.75–1.00 mm) and the content of carbohydrates and OC in these aggregates. The OC and carbohydrates contents of these soils did not account, however, for the variation in aggregate stability of these soils.

### Acknowledgements

This study was supported by the European Union STD Research Project No. TS3 - CT94 - 0281 'Evaluation of the Stability of Soil Organic Matter (SOM) Pools in African Ecosystems (Ethiopia and Nigeria) for the Improvement of a Sustainable Soil Management.' One of the authors (J.S.C.M) acknowledges the financial support from the Abdus Salam International Centre for Theoretical Physics (ICTP), Trieste (ITALY) which enabled him to visit Italy during some aspects of this study.

### References

- Adamu J, Mbagwu JSC & Piccolo A (1997) Carbon, nitrogen and phosphorus distributions in aggregates of forest and cultivated soils in central plateau, Nigeria. In: Drozd J, Gonet SS, Senesi N & Weber J (Eds) The Role of Humic Substances in the Ecosystem and in Environment Protection (pp 257–261). IHSS-Polish Society of Humic Substances, Wroclaw, Poland
- Anderson JM & Ingram JSI (Eds) (1993) Tropical Soil Biology and Fertility Handbook of Methods (2nd ed.). CAB International, Wallington, UK
- Angers DA & Mehuys GR (1989) Effects of cropping on carbohydrate content and waterstable aggregation of a clay soil. Can. J. Soil Sci. 69: 373–380
- Angers DA & Mehuys GR (1990) Barley and alfalfa cropping effects on carbohydrate contents of a soil and its size fractions. Soil Biol. Biochem. 22: 285–288

- Angers DA & N' Dayegamiye A (1991) Effects of manure application on carbon, nitrogen and carbohydrate contents of a silt loam and its particle-size fractions. Biol. Fertil. Soils. 11: 79–82
- Angers DA, Samson N & Légère A (1993) Changes in water-stable aggregation induced by rotation and tillage in a soil under barley production. Can. J. Soil Sci. 73: 51–59
- Arrouays D & Pelissier P (1994) Changes in carbon storage in temperate humic clay soils after forest clearing and continuous corn cropping in France. Plant and Soil 160: 215–223
- Arshad MA, Schnitzer M, Angers DA & Ripmeester JA (1990) Effects of till versus no-till on the quality of soil organic matter. Soil Biol. Biochem. 22: 595–599
- Baldock JA, Kay BD & Schnitzer M (1987) Influence of cropping treatments on the monosaccharide content of the hydrolysates of a soil and its aggregate fractions. Can. J. Soil Sci. 67: 489–499
- Beare MH, Hendrix PF & Coleman DC (1994) Water-stable aggregates and organic matter fractions in conventional- and no- tillage soils. Soil Sci. Soc. Am. J. 58: 777–786
- Benzing-Purdie LM (1980) Organic matter and carbohydrate distribution in an orthic gleysol. Soil Biol. Biochem. 12: 567–571
- Cambardella CA & Elliot ET (1992) Particulate soil organic matter changes across a grassland cultivation sequence. Soil Sci. Soc. Am. J. 56: 777–783
- Cambardella CA & Elliot ET (1993) Carbon and nitrogen distribution in aggregates from cultivated and native grassland soils. Soil Sci. Soc. Am. J. 57: 1071–1076
- Chan KY (1997) Consequences of changes in particulate organic carbon in Vertisols under pasture and cropping. Soil Sci. Soc. Am. J. 61: 1376–1382
- Caron J, Kay BD & Stone JA (1992) Improvement of structural stability of a clay loam with drying. Soil Sci. Soc. Am. J. 56: 1583–1590
- Chesire MV, Christensen BT & Sorensen LH (1990) Labelled and native sugars in particle-size fractions from soils incubated with <sup>14</sup>C straw for 6 and 18 years. J. Soil Sci. 41: 29–39
- Dalal RC & Mayer RJ (1986) Long-term trends in fertility of soils under continuous cultivation and cereal cropping in southern Queensland: I. Total organic carbon and rate of loss from the soil profile. Aust. J. Soil Sci. Res. 24: 281–282
- Dalal RC & Mayer RJ (1988) Cultivation effects on carbohydrate contents of soil and soil fractions. Soil Sci. Soc. Am. J. 52: 1361–1365
- Davidson EA & Ackerman IL (1993) Changes in soil carbon inventories following cultivation of previously untilled soils. Biogeochem. 20: 161–193
- Degens B & Sparling G (1996) Changes in aggregation do not correspond with changes in labile organic C fractions in soil amended with <sup>14</sup>C-glucose. Soil Biol. Biochem. 28: 453– 462
- Dinel H, Righi D, Hardy M & Jambu P (1997) Neutral lipids and structural stability of physically degraded soils. Agrochimica XLI: 97–108
- Dutarte P, Bartoli F, Andreax F, Portal JM & Ange A (1993) Influence of content and nature of organic matter on the structure of some sandy soils from West Africa. Geoderma 56: 459–478
- Elliot ET (1986) Aggregate structure and carbon, nitrogen and phosphorus in native and cultivated soils. Soil Sci. Soc. Am. J. 50: 627–633
- Flanzluebbers AJ & Arshad MA (1997) Soil microbial biomass and mineralizable carbon of water-stable aggregates. Soil Sci. Soc. Am. J. 61: 1090–1097
- Gee GW & Bauder JW (1986) Particle-size analysis. In: Klute A (Ed) Methods of soil analysis, Part 1. 2nd Edition, ASA-SSSA, Madison, WI

- Gupta VVSR & Germida JJ (1988) Distribution of microbial biomass and its activity in different soil aggregate size classes as affected by cultivation. Soil Biol. Biochem. 20: 777–786
- Haynes RJ & Swift RS (1990) Stability of soil aggregates in relation to organic constituents and soil water content. J. Soil Sci. 41: 73–83
- Haynes RJ & Francis GS (1993) Changes in microbial biomass C, soil carbohydrate composition and aggregate stability induced by growth of selected crop and forage species under field conditions. J. Soil Sci. 44: 665–675
- Hu S, Coleman DC, Beare M.H. & Hendrix PF (1995) Soil carbohydrates in aggrading and degrading agroecosystems: influences of fungi and aggregates. Agric. Ecosyst. & Environ. 54: 77–88
- Insam H (1996) Microorganism and humus in soils. In: Piccolo A (Ed) Humic Substances in Terrestrial Ecosystems (pp 265–292). Elsevier, Amsterdam
- Kemper DW & Rosenau RC (1986) Aggregate stability and size distribution. In: Klute A (Ed) Methods of Soil Analysis, Part 1 (pp 425–442). ASA and SSSA, Madison, WI
- Lal R (1986) Conversion of tropical rainforest: potential and ecological consequences. Adv. Agron. 39: 173–263
- Martens DA & Frankenberger WT (1993) Soil saccharide extraction and detection. Plant and Soil 149: 145–147
- Mbagwu JSC & Piccolo A (1998) Water-dispersible clay in aggregates of forest and cultivated soils in southern Nigeria in relation to organic matter constituents. In: Bergström L & Kirchman H (Eds) Carbon and Nutrient Dynamics in Natural and Agricultural Ecosystems (pp 71–83). CAB International, UK
- Mbagwu JSC, Piccolo A & Spallacci P (1991) Effects of field applications of organic wastes from different sources on chemical, rheological and structural properties of some Italian surface soils. Biores. Technol. 37: 71–78
- Monreal CM & Kodama H (1997) Influence of aggregate architecture and minerals on living habitats and soil organic matter. Can. J. Soil Sci. 77: 367–377
- Monreal CM, Schnitzer M, Schulten H-R, Campbell CA & Anderson DW (1995) Soil organic structures in macro and microaggregates. Soil Biol. Biochem. 27: 845–853
- Monreal CM, Schulten H-R & Kodama H (1997) Age, turnover and molecular diversity of soil organic matter in aggregates of a Gleysol. Can. J. Soil Sci. 77: 379–388
- Nelson DW & Sommers LE (1982) Total carbon, organic carbon, and organic matter. In: Page AL, Miller RH & Keeney DR (Eds) Methods of Soil Analysis, Part 2 (pp 539–579). ASA-SSSA, Madison, WI
- Parfitt RL, Theng BKG, Whitton JS & Shepherd TG (1997) Effects of clay minerals and land use on organic matter pools. Geoderma 75: 1–12
- Piccolo A (1996) Humus and soil conservation. In: Piccolo A (Ed) Humic Substances in Terrestrial Ecosystems (pp 225–264). Elsevier, Amsterdam
- Piccolo A & Mbagwu JSC (1990) Effects of different organic amendments on soil aggregates stability and molecular sizes of humic substances. Plant and Soil 123: 27–37
- Piccolo A & Mbagwu JSC (1999) Role of hydrophobic components of soil organic matter on soil aggregate stability. Soil Sci. Soc. Am. J. 63: 1801–1810
- Piccolo A, Zena A & Conte P (1996) A comparison of acid hydrolyses for the determination of carbohydrates in soils. Commun. Soil Sci. Plant Anal. 27: 2909–2915
- Roberson EB, Sarig S & Firestone MK (1991) Cover crop management of polysaccharidemediated aggregation in an orchard soil. Soil Sci. Soc Am. J. 55: 734–738
- Salinas-Garcia JR, Hons FM & Matocha JE (1997) Long-term effects of tillage and fertilization on soil organic matter dynamics. Soil Sci. Soc. Am. J. 61: 152–159

- Smith WN, Rochette P, Monreal C, Desjardins RL, Pattey E & Jaques A (1997) The rate of carbon change in agricultural soils in Canada at the landscape level. Can. J. Soil Sci. 77: 219–229
- Stemmer M, Gerzabek MH & Kandeler E (1997) Soil organic matter and enzyme activity in particle size fractions of soils obtained by low-energy sonication. Soil Biology and Biochem. 30: 9–17
- Stevenson FJ (1994) Humus Chemistry: Genesis, Composition, Reactions (2nd ed.). Wiley Inter Science, New York
- Tisdall JM & Oades JM (1982) Organic matter and water-stable aggregates in soils. J. Soil Sci. 33: 141–163
- Webber LR (1965) Soil carbohydrates and aggregation in crop sequences. Soil Sci. Soc. Amer. Proc. 29: 39–42

tertaile curve charge attraction second in

22