

STRATIFICATION AND LIVESTOCK POPULATION CENSUS FOR ENUGU URBAN, NIGERIA: A PILOT SURVEY

IKPEZE, Obiora Osegboka

Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

ABSTRACT

Stratification and livestock population census for Enugu Urban, Nigeria, between February and April 2005 is described. Based on ground reconnaissance, six stratification zones identified for Enugu Urban (180 km²) were, unplanned Village set-up (9.69 km²), High-density built-up areas (20.25 km²), Medium-density built-up areas (7.90 km²), Low-density built-up areas (9.50 km²), Commercial areas (26.44 km²) and Undeveloped lands/Farms (106.93 km²). About 46.63 km² or 25.90 % of the stratified Enugu Urban was principal suburbs. Estimated livestock population was achieved with 91.70 % of 2927 households in 21 sample blocks of about 8.34 Km² or 17.88 % of the principal suburbs. Livestock population and average population density for stratified Enugu Urban were 32309 (179) for goats, 17027 (95) for sheep, 3765 (21) for pigs, 16152 (90) for dogs, 4338 (24) for cats, 108354 (602) for chickens, 28985 (161) for turkeys and 17160 (95) for ducks. The results of this study may be useful in the formulation of Veterinary, Livestock, Public and Environmental Health Policies, as well as for Livestock Diseases Surveillances, Research Communications and Bioinformatics. The model for this survey could also be adapted for other urban cities in Nigeria and the developing countries of the world where there are no reliable livestock population statistics.

Keywords: Urban stratification, livestock population, diseases surveillances, Bioinformatics.

INTRODUCTION

The civil service, railways and coal mining activities were among the attractions for early immigrant workers to Enugu. Many settlements, which now constitute the principal suburbs, had rapidly sprung up within and at the outskirts of the Coal City. Today, Enugu Urban is a heterogeneous community characterized by social stratifications.

Livestock (goat, sheep, pig, dog, cat, chicken, turkey, and duck) keeping in Enugu was a natural consequence of human settlement. There is now noticeable increase in the number of livestock on free range within the City. Many domestic animals, which strayed about the city, had constituted public menace. Moreover, the situation whereby domestic animals were allowed to scavenge on street garbage, refuse dumps, sewage effluents, slaughter premises, markets, shallow streams, open parks, farmlands, and school premises etc., constituted potential hazards to public and environmental health. *Toxocara canis*, one of the commonest parasites of dogs in many parts of the world (Woodruff, 1975; Stewart *et al.*, 1979), is the most important cause of Visceral Larval Migrants (VLM) in man (Beaver, 1956). Chiejina and Ekwe (1986) had reported on the environmental contamination associated with *Toxocara* eggs in dog faeces in Enugu and Nsukka. Dada and Belino (1979), Onadeko and Ladipo (1989) also described the public health significance of Ascariasis, Trichuriasis and helminthes' ova in dog faeces from Nigerian urban towns. Rabies is also transmitted to man through the bite of a rabid dog. 'Street rabies virus' (SRV) had been demonstrated in the saliva of dogs (Vaughn *et al.*, 1965). Other human infections like trichinosis and hydatid diseases

may be contracted from infested pork and beef, respectively.

The aim of this pilot study was to establish stratification zones in Enugu Urban for an effective conduct of livestock population census, test the instruments of urban livestock population census, assess the reliability and usability of the demarcation maps and ascertain the time that may be required to conduct livestock population census for Enugu Urban. Since there is a dearth of information on livestock population figures in Enugu and other urban cities of Nigeria, data from this pilot survey and related studies may be useful in the formulation of Veterinary, Livestock, Public and Environmental Health Policies, as well as for Livestock Diseases Surveillances, Research Communications and Bioinformatics.

MATERIALS AND METHODS

Ground Survey: Street map of Enugu Urban obtained from the Department of Lands, Surveys and Urban Planning, Enugu was used for this survey. Based on ground reconnaissance, six stratification zones were established. The extent of each stratum was demarcated on the stratification map of Enugu while the areas (Km²) were determined by the use of a squared graph paper. Twenty-one sample blocks were randomly selected from the principal suburbs of Enugu (excluding commercial and farmlands which had no permanent livestock). The extent of the 21 sample blocks were established physically on the ground both by pacing (a pace \approx 1 metre) and with the aid of a pedometer strapped to the hip belt. Guides to the extent of the strata and sample blocks were produced (Figure 1).

Table 1: Stratification zones and sample blocks in Enugu Urban*

Strata	Stratification Zones			No.	Area (Km ²)	Bounding streets	Principal Suburbs (where sample blocks were located)				
	Text Abbreviation	Area (Km ²)	% of Urban Enugu								
Unplanned Village Set-up	'Village'	9.69	5.38	1	0.76	Entire Agu-Abor	Agu Abor				
				2	0.56	Half of Ugbo Odogwu	Ugbo Odogwu				
				3	0.64	Entire Ugwu Aaron	Ugwu Aaron				
				4	0.70	Entire Ugwu Alfred	Ugwu Alfred				
High density areas	'Township'	20.25	11.25	5	0.24	Nkpor – Atani – Ikem Streets	Abakpa				
				6	0.18	Inyi – Obioma – Umunwakum Streets.	Achalla Layout				
				7	0.40	Agbani Road – Umueze Street	Awkunanaw				
				8	0.14	Colliery Quarters	Coal Camp				
				9	0.36	Chima Avenue – Kano Street	New Haven				
				10	0.46	Agbani Road – Gold Smith Ave.	Ogbete				
				11	0.26	Edinburgh Road – Kenyatta St.	Ogui New Layout				
				12	0.22	Zik's Ave – Christ Church – Agbani Rd.	Uwani				
				Medium density areas	'Low-cost'	7.19	4.00	13	0.38	Ichida Street – Osina Street	Federal Housing
								14	0.28	Ogui Rd – 1 st – 4 th Avenue	Artisan Quarters
								15	0.86	Imoke Street – Ekulu St – Nwodo Ave.	Trans-Ekulu Housing
				Low density areas	'GRA'	9.50	5.27	16	0.60	NCO Blocks	Army Barracks
17	0.16	Abakaliki Rd – Army Barracks	Army Officers' Qtr.								
18	0.30	Rangers' Ave – Ezzikwo Street	Modern Residential								
19	0.38	Abakaliki Lane – Charles Street	"GRA"								
20	0.26	Riverside Estate – Air Force Base	Thinkers' Corner								
21	0.20	ESUTH – WTC - UNEC	Campus Residential								
Commercial areas	'Industrial'	26.44	14.70	Livestock census was not carried out in 'Industrial' & 'Open' (Uninhabited Urban Enugu) where no permanent livestock existed							
Undeveloped lands/Farms	'Open'	106.93	59.40								
Stratified Urban Enugu		180.00	100.00								

*Total area of sample blocks in 'Village' = 2.66 Km² (27.45 % of 'Village'), Total area of sample blocks in 'Township' = 2.26 Km² (11.16 % of 'Township'); Total area of sample blocks in 'Low-cost' = 2.12 Km² (29.48 % of 'Low-cost'); Total area of sample blocks in 'GRA' = 1.30 Km² (13.68 % of 'GRA'); Total area of sample blocks = 8.34 Km² (17.88 % of inhabited Enugu); Area of inhabited Enugu = 46.63 Km² (25.90 % of stratified Enugu); Area of uninhabited Enugu = 133.37 Km² (74.10 % of stratified Enugu)




Figure 1: Map of Enugu urban showing the principal suburb and 21 sampled blocks

Livestock Enumeration: Three pairs of livestock assistants and a livestock superintendent served as enumerators and field supervisor respectively, while the researcher coordinated all activities. Field workers, provided with logbooks, were guided through the extent of the sample blocks. There were seven sample blocks per pair of enumerators.

Enumeration in each sample block required only two days (i.e., Saturday and Sunday) of a week, so that the entire fieldwork lasted for 14 working days spread in-between 7 weeks from February to April 2005. Data on every sample size (household) and species (livestock) enumerated were entered differently in the logbooks.

Collation and Analysis of Data: All data entries in the logbooks were collated and analyzed to obtain the estimated populations and population densities of each species of livestock in Enugu. Absolute numbers of animals in a stratum were obtained after multiplying the totals for the sample blocks in that stratum by 'a factor', which was the percentage area of that stratum represented by the sample blocks. The average livestock population density for Enugu (including commercial and farm lands) was obtained by dividing the 'total' for that livestock species by the area of stratified Enugu Urban.

RESULTS AND DISCUSSION

The stratification of Enugu is illustrated in Figure 1. Six stratification zones (Strata) described for Enugu were unplanned zones (Strata) described for Enugu were unplanned zones ('Village'), High-density areas ('Township'), Medium density areas ('Low-cost'), Low-density areas ('GRA'), Commercial areas ('Industrial') and Undeveloped lands/Farmlands

Table 2: Livestock species enumerated from 21 sample blocks in stratified Enugu

Strata	Sample Blocks	Principal Suburbs	NR*	Households				Livestock						
				NL	LA	TH	Goat	Sheep	Pig	Dog	Cat	Chicken	Turkey	Duck
Village	1	Agu Abor	5	4	113	122	93	35	15	40	8	211	30	50
	2	Ugbo												
	3	Odogwu	3	3	121	127	58	24	25	35	5	521	51	42
	4	Ugwu												
Township	5	Aaron	4	5	104	113	76	49	19	37	16	152	23	21
	6	Alfred	9	3	108	120	68	61	29	29	9	273	18	28
	7	Abakpa	5	1	182	188	36	18	40	54	5	610	51	128
	8	Achalla												
	9	Layout	4	3	156	163	53	42	18	45	7	421	20	92
	10	Awkunanaw	12	1	170	183	116	73	37	28	8	720	210	281
	11	Colliery	3	2	70	75	75	38	-	19	5	241	53	51
	12	New Haven	7	4	149	160	43	14	-	23	17	308	61	30
	13	Ogbete	2	-	165	167	143	70	16	76	21	652	43	69
	14	Ogui New												
	15	Layout	2	15	80	97	204	96	-	35	10	446	52	45
Low-cost	16	Uwani	11	2	210	223	127	74	10	23	15	390	95	109
	17	Fed.												
	18	Housing	8	20	121	149	71	33	-	41	8	271	110	15
	19	Artisan Qtr	9	14	110	133	144	81	-	59	7	188	232	25
GRA	20	Trans Ekulu												
	21	Hous.	2	12	124	138	77	35	-	65	18	171	181	39
	22	Army Barracks	10	11	80	101	120	57	-	14	13	280	25	43
	23	Army Offs' Qtr	3	2	78	83	28	13	-	16	9	75	56	13
	24	Mod. Residential	3	1	125	129	58	29	-	46	16	63	27	9
	25	"GRA"	9	5	167	181	71	33	-	88	21	81	46	18
TOTAL	26	Thinkers' Cn. Campus Residential	5	-	130	135	45	28	-	71	11	121	37	12
TOTAL			15	3	122	140	30	12	-	44	13	208	50	-
TOTAL			131	111	2685	2927	1736	915	209	888	242	6403	1471	1120

*NR = No response from Household (4.5 %); NL = No livestock owned by household (3.8 %); LA = Livestock owned by household (91.70 %); TH = Total number of households visited (2927)

Table 3: Estimated livestock population and population density of stratified Enugu

Species	Total Sample Blocks				Estimated Livestock Population					Population Density
	'Village'	'Township'	'Low-cost'	'GRA'	Village (27.45)*	Township (11.16)	Low-cost (29.48)	GRA (13.68)	Total	
Goats	295	797	412	232	8097	8894	12145	3173	32309	179
Sheep	169	425	206	115	4639	4743	6072	1573	17027	95
Pigs	88	121	-	-	2415	1350	-	-	3765	21
Dogs	141	303	179	265	3870	3381	5276	3625	16152	90
Cats	38	88	46	70	1043	982	1356	957	4338	24
Chickens	1157	3788	910	548	317	42274	26826	7496	108354	602
Turkeys	122	585	548	216	3348	6528	16155	2954	28985	161
Ducks	141	805	122	52	3870	8983	3596	711	17160	95

* Absolute numbers of livestock in a stratum were obtained after multiplying the totals for all the sample blocks in each stratum by 'a factor (in brackets)', i.e., the percentage area of that stratum represented by the sample blocks. The average population density was obtained by dividing the 'Total' column by 180 Km², which was the area of stratified Enugu.

('Open'). Iyioku, Nyaba, Asata, Ayo, Ekulu and Idaw rivers and Awa stream drain the Coal City, which is traversed by the Onitsha-Enugu-Port Harcourt Expressway.

Principal Suburbs and twenty-one (21) sample blocks are represented in Figure 1 while Table 1 shows the extent of the six (6) stratification zones (strata) and the twenty-one (21) sample blocks surveyed in Enugu Urban (180 km²). Unplanned village set-ups (9.69 km² or 5.38% of stratified Enugu Urban) are nearly as old as the city and are located mostly on the outskirts of the city. 'Village' included Agu Abor, Agu Ugwu, Gabon, Obinagu, Onu Ogba

Nike, Ugbo Odogwu, Ugwu Aaron, Ugwuaji and Ugwu Alfred. High-density townships (20.25 km² or 11.25 % of stratified Enugu) included Abakpa, Achalla layout, Asata, Awkunanaw, Idaw-river layout, Mainland, Mary land, New Haven, Ogbete (Coal camp), Ogui, Ogui New layout, Udi Siding (PWD and P and T Quarters), Ugwu Ago Nike, and Uwani. Medium-density areas (7.19 km² or 4.0 % of stratified Enugu) included the Federal low-cost houses, Trans Ekulu layout, Army barracks, and Railway quarters. Low-density 'GRA' category (9.50 km² or 5.27 % of stratified Enugu) included the Abakaliki Road GRA, Independence Layout, Government House, Modern and Campus

residential areas, Senior Army Officers quarters, and Thinkers corner.

Commercial areas (26.44 km² or 14.70 % of stratified Enugu) included the Air Force base, Emene industrial, Markets, Nike Lake Hotels, Ogui commercial, Secretariat, and WTC/Queens/ESUT/IMT areas. Undeveloped areas (106.93 km² or 59.40 % of stratified Enugu) were made up of farmlands and undeveloped plots at Akwuke, Amaoji, New GRA, Nkwubo Nike, Obinagu, Onu Ogba Nike, Ubaha Nike, and Ugwuaji, where permanent livestock were not observed. About 46.63 km² (25.90 %) of Enugu was regarded as inhabited and from which the 21 sample blocks (8.34 km² and about 17.88 % of inhabited Enugu) were located. Permanent livestock were observed here.

Data on household and livestock enumeration are presented in Table 2. 2927 households were visited during the exercise. Occupants of 131 (4.5 %) were either absent or did not respond. Occupants of 111 (3.8 %) had no livestock while those in 2685 (91.70 %) had one type of livestock species or the other.

Table 3 showed the estimated livestock population census figures for stratified Enugu (including 'Industrial' and 'Open' areas). The average population densities of the different livestock species (number per km²) were estimated to be in the order of chickens (602) > goats (179) > turkeys (161) > sheep and ducks (95) > dogs (90) > cats (24) > pigs (21).

Large herds of nomadic cattle and many commercial poultry farms encountered during the survey, no doubt contributed to urban environmental contamination and pollutions but were not included in this pilot study, which was limited to small animals and ruminants that were usually kept in the 'back yard', allowed on free range and which frequently strayed within the urban city.

This study effectively tested the instruments of urban livestock population census, assessed the reliability and usability of the demarcation maps and ascertained the time that may be required to conduct livestock population census for Enugu Urban. Since the percentage of non-cooperation was ≤ 5%, the result may not contain any great 'unknown' within its figures. The study was also

productive of results without waste of labour. The model for this pilot survey could also be adapted for use in other cities of the country and in the developing countries of the world where livestock population statistics are either unavailable or unreliable.

ACKNOWLEDGEMENT

I thank Mr. Peter Okafor (a Cartographer with the Ministry of Lands, Survey and Urban Planning, Awka), all the livestock assistants and their superintendent who assisted in the accomplishment this arduous task. I am grateful to Dr (Mrs.) N. N. Ezumah of Gender and Developmental Research Group (GADREG) - an Enugu based Non-Governmental Organization (NGO) - for financing this pilot study.

REFERENCES

- BEAVER, P. C. (1956). Larva migrans. *Experimental Parasitology*, 5: 587 – 621.
- CHIEJINA, S. N. and EKWE, T. O. (1986). Canine toxocariasis and the associated environmental contamination of urban areas in eastern Nigeria. *Veterinary Parasitology*, 22: 157 – 161.
- DADA, B. D. J. and BELINO, E. O. (1979). Prevalence and public health significance of helminth ova in dog feces deposited on streets in Zaria. *Annals of Tropical Medicine and Parasitology*, 73: 495.
- ONADEKO, M. O. and LADIPO, A. O. (1989). Parasitic infestation in rural communities: A focus for primary health care in Nigeria. *African Journal of Medical Science*, 18: 289 – 294.
- STEWART, G. L., REDDINGTON, J. J. and SMITH, W. G. (1979). Intestinal helminth parasites of dog from Terrant County, Texas. *Southwestern Veterinarian*, 32: 29 – 32.
- VAUGHN, J. B., GERHARDT, P and NEWELL, K. W. (1965). Excretion of street rabies virus in the saliva of dogs. *Journal of American Veterinary Association*, 193 (5): 113.
- WOODRUFF, W. A. (1975). *Toxocara canis* and other nematodes transmitted from dogs to man. *British Veterinary Journal*, 131: 627 – 632.