

***A  
N  
I  
M  
A  
L***



***R  
E  
S  
E  
A  
R  
C  
H***



***I  
N  
T  
E  
R  
N  
A  
T  
I  
O  
N  
A  
L***



**An International Journal Publishing Original Research Involving  
the Use of Animals and Animal Products**

**ISSN: 159-3115**

**Website: [zoo-umn.org](http://zoo-umn.org)**

## MACROBENTHIC FAUNA OF A HUMID TROPICAL WATER RESERVOIR, ABIA STATE, NIGERIA

<sup>1</sup>AVOAJA, Diana Akudo, <sup>2</sup>INYANG, Nicholas Mathias and <sup>3</sup>MGBENKA, Bernard Obialo

<sup>1</sup>Department of Biological Sciences, Michael Okpara University of Agriculture, Umudike Umuahia, Abia State, Nigeria

<sup>2</sup>Fisheries and Hydrobiology Research Unit, Department of Zoology, University of Nigeria, Nsukka, Enugu State, Nigeria

**Corresponding Author:** Avoaja, D. A., Department of Biological Sciences, Michael Okpara University of Agriculture, Umudike, PMB 7267, Umuahia, Abia State, Nigeria. Phone: 23248063573664

### ABSTRACT

*The macrobenthic fauna of a humid tropical zone water reservoir was investigated from January to December 2002. Benthos was obtained using an improvised Surber Stream Bottom Sampler. Three phyla of macrobenthos (Arthropoda, Mollusca and Annelida) prevail in the reservoir, with quantitative variations. A total of 1,279 macrobenthic animals were recorded. The Arthropoda was represented by larvae of five genera of insects, with a total of 644 (50.4 %), while Mollusca was represented by two genera with a total of 165 (12.9 %), and Annelida by three genera, with a total of 470 (36.7 %) of the benthos. There was significant difference ( $P < 0.05$ ) among the phyla populations of the benthos.*

**Keywords:** Macrobenthic, fauna, Humid, Tropics, Water reservoir

### INTRODUCTION

The water reservoir is located in the National Root Crops Research. Institute (NRCRI), Umudike. This Institute is situated 8 kilometers south east of Umuahia, in Abia State. The reservoir was impounded in 1965 from a tributary of the river Qua-Iboe, which passes through the Institute, for irrigation of farms during dry seasons. As a result of the long existence of the reservoir, and erosion of the banks, it has become a large body of water, with an estimated surface area of 15.5 km<sup>2</sup>. (Avoaja, 2005) The climate of Umudike is typical of the humid tropics, with fairly even and uniform temperatures throughout the seasons of the year (Iloeje, 1980). The rainy season usually starts from March and ends in October, with relatively constant temperature, frequent rains, and high humidity (Iloeje, 1980).

Both qualitative and quantitative studies on the benthos of lentic and lotic waters abound in literature (Williams and Hynes, 1971). Poor taxonomic knowledge of African freshwater faunas in general, allows few groups to be identified beyond the generic level, thus limiting detailed analysis of benthic communities. (Williams and Hynes, 1971) According to Vareschi and Vareschi (1984), the benthic fauna of lake Nakuru (Kenya) was remarkably poor in species, which consisted two chironomid species. Lake Chad in contrast had 47 chironomids (Dejoux, 1968). Nematodes were found only very occasionally and the undersides of stones along the western shores of the lake were populated by the coleopteran, *Helochaetes species (Hydrophilidae)*. Characteristic benthic species of other lakes e.g. Oligochaetes, Chaoboridae, Ostracods or Molluscs were completely absent in lake Nakuru (Vareschi and Vareschi, 1984). In a tropical flood river basin, the common invertebrate taxa were crustaceans, insect

and gastropod mollusks (Ezenwaji, 1982; Okafor, 1990; Eyo and Ekwonye, 1995). Banerjee and Banerjee (2005) identified five types of macrobenthic fauna (Oligochaeta, Polychaeta, Crustacea, Gastropoda, and Bivalva) in three Indian estuaries.

Imbevbor and Bakare (1970) reported that most benthos and macroinvertebrates fed on the debris that settled at the bottom of water, and in turn served as food for a wide range of fish. Similar observations have been reported (Moss *et al.*, 1987; Eyo and Ekwonye, 1995). The identification of the benthos of this reservoir would give an insight into the natural fish food of the water body. The present paper embodies a study of macrobenthic fauna of the reservoir in Umudike, which adds to the knowledge of distributions and abundance of macrobenthic fauna in tropical freshwater ecosystem.

### MATERIALS AND METHODS

Investigation was carried out in the reservoir from January to December 2002. The reservoir was divided into six sampling stations by a transect method. The six sampling stations were approximately two kilometers apart. That gave a stratified sampling station in which sub-divisions of the stations were sampled using a random sampling technique.

Benthos were sampled using an improvised Surber Stream Bottom Sampler, (a net 0.5 cm mesh, with an area of 900 sq. cm, on a metal frame). A sampling site in each of the stations was selected. The net was dipped inside the water (about 0.5 m depth) collecting mud with the edge of the square metal, into the net. The procedure was repeated five times, and each sample was put into different jars. The jars were labeled indicating the number of replicates, and then carried to the laboratory.

**Table 1: Monthly macrobenthos collections from the reservoir**

Months	Annelida			Mollusca			Arthropoda				Total	No. /M <sup>2</sup>
	<i>Nais batata</i>	<i>Nais simplex</i>	<i>Glossiphonia sp.</i>	<i>Lymnaea sp.</i>	<i>Physella sp.</i>	<i>Chironomus sp larvae</i>	<i>Dysticus sp. larvae</i>	<i>Copelatus sp. larvae</i>	<i>Letes sp larvae</i>	<i>Ecdyonurus sp. larvae</i>		
Jan.	7	8	14	5	5	12	8	8	13	10	90	18
Feb.	11	8	10	9	5	12	11	10	11	12	99	19
Mar.	12	14	16	11	4	7	13	8	9	10	104	21
Apr.	13	15	17	10	5	10	8	11	13	14	116	23
May	11	15	14	14	8	8	8	6	11	12	107	21
Jun.	11	11	27	7	7	8	4	9	14	13	111	23
July	13	13	22	6	5	3	8	14	11	14	109	23
Aug.	7	6	32	8	7	13	12	9	13	15	122	23
Sept.	10	12	35	9	7	6	7	7	9	8	110	22
Oct.	7	6	19	6	5	15	17	9	15	14	113	23
Nov.	8	7	13	5	5	13	15	13	13	14	106	21
Dec.	8	7	11	8	4	10	11	10	11	12	92	20
Year	118	122	130	98	67	117	122	114	143	148		
Total Phyla		470			165			644				
Total % age		36.7%			12.9%			50.4%				

**Table 2: Benthos distribution in stations in the water reservoir**

Phylum	Class	Order	Family	Species	Stations					
					1	2	3	4	5	6
Annelida	Oligochaetae	Naidida	Naididae	<i>Nais Simplex</i>	-	-	+	+	+	+
				<i>Nais batata</i>	-	-	+	+	+	+
				<i>Glossiphonia complonata</i>	+	+	+	+	+	+
Arthropoda	Insecta	Diptera	Chironomidae	<i>Chironomus sp. (larvae)</i>	+	+	+	+	+	+
Arthropoda	Insecta	Coleoptera	Dytiscidae	<i>Dysticus marginalis (larvae)</i>	-	+	+	+	+	-
"	"	"	"	<i>Copelatus sp. (larvae)</i>	-	+	+	-	-	-
"	"	Odonata	Lestidae	<i>Lestes sp. (larvae)</i>	+	+	+	+	+	+
"	"	"	"	<i>Ecdyonurus sp. (larvae)</i>	+	+	+	+	+	+
Mollusca	Gastropoda	-	Lymnaeiidae	<i>Lymnaea natalensis</i>	-	-	+	+	+	-
"	"	-	Physidae	<i>Physella sp.</i>	-	-	+	+	+	-

Key: + Present, - Absent

In the laboratory, the contents of each jar were poured into a white enamel pan. The material from each enamel pan was then poured into a fine sieve (a no. 40 U.S. series sieve) to remove water, sand and debris. The organisms in the sieve were sorted out manually, and separated into taxonomic groupings for final identification and enumeration. The procedure was carried out for each of the six stations. The number of benthos per square meter was computed using Lind (1979) thus: number of benthos per meter = Total no per month / Total area sampled

(m<sup>2</sup>) where, Total area sampled = area of net x No of replicates.

The animals recovered were put into specimen bottles, labelled, and preserved in 4 % formaldehyde solution for further examination. These specimens were identified using keys (Edmondson, 1966).

**Statistical Analysis:** SPSS version 10 employing simple percentiles and graphs were used in analyzing

the population of macrobenthic fauna of the reservoir.

## RESULTS AND DISCUSSION

The benthic fauna was represented by three animal phyla, Annelida, Mollusca, and Arthropoda. The annelids were Oligochaetes, represented by *Nais simplex* and *Nais batata* and Hirudinea represented by *Glossiphonia complonata*. The Molluscs were the Gastropods, represented by *Lymnaea natalensis* and *Physella* sp. The Arthropods were mainly larvae of insects, represented by the larvae of *Chironomus* sp.; *Dysticus* sp.; *Copelatus* sp.; *Lestes* sp. and *Ecdyonurus* sp. (Table 1). A total of 1,279 benthic specimens were collected during the year. The minimum and maximum benthoses per square metre were recorded. 18 and 23 respectively. The record showed that in terms of number, Arthropoda was the highest with 644 (50.4 %), followed by the Annelida, 470 (36.7 %), and the least was the Mollusca, 165 (12.9 %). Table 2 shows the benthic distribution in the stations.

The Gastropods were found in stations 3, 4 and 5. The insect larvae were found almost in all the stations but stations 1 and 6 recorded no *Dysticus* sp. larvae, and stations 1, 4, 5 and 6 recorded no *Copelatus* larvae. There were significant differences ( $P < 0.05$ ) between the phyla of the macrobenthos of the reservoir.

Benthic organisms play several important roles in the aquatic community. They are involved in the mineralization and recycling of organic matter produced in the open water or brought from external sources (Moss *et al*; 1987). They are important second and third links in the trophic sequence of aquatic communities (Moss *et al*; 1987). Many benthic insect larval forms are major food sources for fishes (Lind, 1979). These insect larvae identified in this study were probably food for both the young and adult fishes in the reservoir. Few animals which have the ability to exploit the benthic mud region of lakes and reservoirs, because of low oxygen tension arising from decay of organic matter, survive. Therefore, the degree of specialization required from the organisms in order to survive in the mud region means that the number of species will be limited (Brown, 1971). This probably, accounted for the few molluscs and Annelids recovered in this study.

## REFERENCES

- AVOAJA, D.A. (2005). *The Ecology and fish fauna of the Umudike water reservoir in Abia State Nigeria. PhD Thesis*, University of Nigeria Nsukka. 182 pp.
- BANERJEE, P. and BANERJEE, S. (2005). Macrobenthic fauna in relation to soil and water quality of a portion of Hooghly Estuary from Konagar to Bally, West Bengal. *Environment and Ecology* 23(3): 687-690.
- BROWN, A. L. (1971). *Ecology of fresh water*. Heinemann Educational Books Limited, London. 129 pp
- DEJOUX C. (1968). Contribution a L'etude des insectes aquatiques du Tchad. Catalogue des Chironomidae, haoboridae, Odonates, Trichoptere, Hemiptere, phemeropteres. Cah. ORSTOM Ser. *Hydrobiology*, 11(2): 51 – 78.
- EDMONDSON, W. T. (1966). *Freshwater Biology* 2<sup>nd</sup> Ed. New York, London Sydney. 1248 pp.
- EYO, J. E. and EKWONYE, U. C. (1995). The macroinvertebrate fauna of pool in the floodplain (Fadama) of the Anambra River, Nigeria. *Freshwater Forum*, 5(3): 160 – 162.
- Ezenwaji, H. M. G. (1982). *Aspects of the Biology of Clarias albopunctatus in the Anambra River basin, Nigeria*. M.Sc. Thesis, University of Nigeria Nsukka. 173 pp.
- ILOEJE, N. P. (1980). *A New geography of Nigeria*. Longman Nigerian Limited, Lagos. 200 pp.
- IMEVBORE, A. M. A. and BAKARE, O. (1970). The food and feeding habits of non-cichlid fishes of the River Niger. Pages 49 – 64. In: *The Kanji Reservoir Area. Volume I, Ecology*. University Press, Ibadan.
- LIND, O. T. (1979). *A hand book of common methods in Limnology*, 2<sup>nd</sup> Edition. C.V. Mosby Company, London. 197 pp.
- MOSS, D., ARMILAGE, P. G. and GUNN, R. T. (1987). The use of prediction to assess macro-invertebrate fauna of unpolluted running water sites in Great Britain, using environmental data. *Freshwater Biology*, 17: 41 – 52.
- OKAFOR, F. C. (1990). Distribution of freshwater gastropods in the lower River Niger and Cross River basin of South-eastern Nigeria, with reference to their trematode infections. *Batr. Tropical Landwirtsch Veterinary Medicine*, 28: 207 – 216.
- VARESCHI, E. and VARESCHI, A. (1984). The ecology of Lake Nakuru (Kenya) IV. Biomass and distribution of consumer organisms. *Oecologia*, 61: 70 – 82.
- WILLIAMS, T. R. and HYNES, H. B. N. (1971). A survey of the fauna of streams on mount Elgon, East Africa, with special reference to the Simuliidae (Diptera). *Freshwater Biology*, 1: 227 – 248.