

EVALUATION OF TAMARIND (*Tamarindus indica*) SEED MEAL AS A DIETARY CARBOHYDRATE FOR THE PRODUCTION OF NILE TILAPIA *Oreochromis niloticus* (L)

NWANNA, Lawrence., FAGBENRO, Oyedapo and OLANIPEKUN, Sunday

Department of Fisheries and Wildlife, Federal University of Technology, PMB 704 Akure, Nigeria

Corresponding author: Dr. NWANNA, Lawrence. Department of Fisheries and Wildlife, Federal University of Technology, PMB 704, Akure, Ondo State, Nigeria.

ABSTRACT

A feeding study was conducted to assess the value of Tamarind, Tamarindus indica seed meal as dietary carbohydrate in the diets of Nile Tilapia, Oreochromis niloticus. Tamarind seeds were used to replace maize at 0, 20, 40, 60, 80, 100 % substitution levels for treatments 1 to 6. Growth trial was conducted in outdoor concrete tanks for 56 days. The fishes were fed at 4 % body weight twice daily. There were no significant variations in the mean weight gain, specific growth rate (SGR), food conversion ratio (FCR) and protein efficiency ratio (PER) ($p > 0.05$). The apparent digestibility coefficient (ADC) of protein and energy of the fishes fed diets 1 – 6 were similar ($p > 0.05$). There were no significant differences in the blood total erythrocyte counts, (TEC), pack cell volume (PCV) and red blood cells count (RBC) ($p > 0.05$). Based on the findings, complete replacement of maize with tamarind seed meal in the diets of O. niloticus is recommended.

Keywords: Tamarind, Replacement, Maize, *Oreochromis niloticus*.

INTRODUCTION

Plant protein and carbohydrate feedstuffs, particularly soybean (*Glycine max*) and maize (*Zea mays*), have been extensively used in fish feeds with nutritional, environmental and economic benefits (Tacon, 1993). As the use of soybeans and maize in human food and livestock feed increases, their costs have increased, and the economics of using them in fish feeds may become less favourable. Hence the evaluation of underutilized indigenous plant protein/energy-rich sources becomes imperative and remains a high research priority in Nigeria.

Vast quantities of forest seeds are discarded as wastes in Nigeria. There is a strong economic justification for their use either as protein or carbohydrate/energy supplements in low cost diets for fish. A national feedstuff survey revealed that the seed of tamarind, (*Tamarindus indica*) represents a good source of digestible dietary energy and is desirable in fish feeds because of its low cost and availability (Balogun 1990). Tamarind contains about 18 % crude protein, with amino acid profile comparable to maize, but with a higher methionine + cystine (met + cys) value than maize. There is a dearth of information on the nutritive value of tamarind seed meal as a feedstuff for Nile tilapia, *Oreochromis niloticus*.

Oreochromis niloticus is a fast growing and preferred aquaculture species in Africa and is capable of utilizing plant materials in its diets. The

present study was conducted to evaluate the nutritional and economic feasibility of replacing maize with tamarind seeds in practical diets for *O. niloticus*.

MATERIALS AND METHODS

Diet: The ingredients used in this study were purchased from Pfizer (Livestock) feeds depot, Ibadan Nigeria, while tamarind seeds were obtained from Kainji lake area, New Busa, Nigeria. Tamarind seeds were mechanically de-hulled. The dehulled seeds were soaked in warm water for 24 h to remove the inner coat before sun drying to a constant moisture of < 10 %, and milled into fine powder. Phytin and tannin contents of the tamarind seed meal (TM) were determined according to the methods of Sathe and Salunkhe (1981). The mineral content of TM was determined as described by Harris (1970), while the proximate composition was determined according to the methods of AOAC (1990) (Table 1).

Based on the proximate composition of the feedstuffs, six diets were formulated (Table 2). The control diet (diet 1) contained 30% of maize which was replaced with TM in diets 2, 3, 4, 5 and 6 at 20, 40, 60, 80 and 100% respectively. The feedstuffs were milled, blended, moistened, pelleted, sun-dried at 30°C between 1200 – 1600h for three days and stored in air-tight polythylene bags at ambient

temperature (28°C). Proximate analysis of the diets and the fishes, (moisture, crude protein (N 58. x 6.25), crude lipid, crude fibre and total ash) were conducted in triplicate samples according to AOAC (1990) methods. Gross energy of the diets was determined in triplicate samples by combustion in bomb calorimeter.

Growth Trials: Groups of 20 fingerlings of *O. niloticus* (6.15 ± .03g) having been acclimated for seven days were randomly stocked into 18 outdoor rectangular concrete tanks (1.8 x 1.8 x 1m) containing 800 L of water for growth trials. The mean pH, dissolved and temperature of the tank waters were 6.9 ± 0.2, 5.6 ± 0.4 and 26.8 ± 0.6 respectively. Each of the diets was fed to the fishes in triplicate tanks at 4 % body weight twice daily (0900–1000 and 1500–1600h) for 56 days. Fresh water was used to change the water in the experimental tanks once every two weeks. Total weight of fishes in each tank was taken bi-weekly to monitor growth responses and for feed adjustments. Mean weight gain (MWG) specific growth rate (SGR), protein efficiency ratio (PER) and food conversion ratio (FCR) were estimated from bi-weekly weight data according to the methods of Olivera Novoa *et al.* (1990) as follow: Mean weight gain = final mean weight – initial mean weight; Specific growth rate = $10^2 (\text{Log}_e \text{ final weight} - \text{Log}_e \text{ initial weight}) / \text{culture period (days)}$; Food conversion ratio = Dry weight of feed fed (g)/ fish weight gain; Protein efficiency ratio = fish weight gain/protein fed.

Carcass and Haematological Analyses: At the beginning and end of feeding trials, six tilapia, randomly selected from each treatment group were oven dried at 48 °C for 48 h, blended into fine powder, packed in air-tight polythene bags and stored in a deep freezer (-20°C), prior to carcass analyses. Similarly, before and after the feeding trials three fish from each tank (9 fish/treatment) were removed, anesthetized using 10 mg/l tricaine methane sulfonate (MSS222, Sandoz). Blood samples were withdrawn from the caudal vein with heparinized syringes, and immediately centrifuged at 5,000 rpm for 15 mins to remove red blood cells. The blood parameters, total erythrocyte count (TEC), pack cell volume (PCV) and red blood cells (RBC) were determined according to the methods of Svobodova *et al.* (1991).

Digestibility Studies: Groups of 20 *O. niloticus* fingerlings (7.2 ± 0.8g) having been

acclimated for seven days were randomly stocked into 18 indoor 60 L rectangular glass tanks (75 x 40 x 40 cm) supplied with fresh water (flow rate 1L m⁻¹). Each of the diets was fed to the fishes in triplicate tanks at 4 % body weight twice daily for 14 days. Faeces were collected from each of the tanks eight hours after each feeding. The faeces and unfed diets were pooled separately and their proximate composition determined according to the methods of AOAC (1990). The digestibility of unfed feed and the dried faeces per sample/treatment were determined by AIA method (Halver *et al.* 1993). These samples were ashed and digested by Acid in soluble Ash (AIA). $\text{AIA}\% = 10^2 (\text{weight of Ash} - \text{weight of AIA}) / \text{weight of Ash}$. Apparent digestibility coefficient (ADC) was determined from the formula, $\text{ADC} = 10^2 - (10^2 \times (\text{Af}/\text{At} \times \text{Nt}/\text{Nf}))$ where, Af = AIA in feeds, At = AIA in faeces, Nf = Nutrient in feed and Nt = Nutrient in faeces.

Statistical Analyses: The one way analysis of variance (ANOVA) and Duncan's multiple-range tests (Zar, 1984) were used to compare differences between diet treatment means (p = 0.05).

RESULTS AND DISCUSSION

Table 1 presents the mineral, proximate composition, and anti-nutrients in tamarind seeds; which showed potassium (K), calcium (Ca), and magnesium (Mg), as the most abundant minerals. The protein content is high (16.6%), with low fat and fibre contents.

Table 1: Mineral, anti-nutrient and proximate composition of tamarind seed meal

Items Analyzed	Proximate Composition (g/100g/DM)
Mineral and Anti-nutrient (g/kg)	
Iron (Fe)	0.10
Zinc (Zn)	0.13
Magnesium (Mg)	0.31
Sodium (Na)	0.23
Potassium (K)	0.43
Calcium (Ca)	0.40
Phytin	1.17
Tannin	0.02
Nutrients	
Crude Protein	16.6
Fat	0.35
NFE	69.8
Moisture	4.74
Fibre	5.82
Ash	2.70

Table 2: Ingredient and proximate composition of experimental diets

Items Analyzed	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Diet 6
Ingredients(g/100g/DM)						
Fish meal	15	15	15	15	15	15
Soybean meal	45	45	45	45	45	45
Maize	30	24	18	12	6	0
Tamarind	0	6	12	18	24	30
Fish oil	6	6	6	6	6	6
Vitamin-Mineral Premix ¹	2	2	2	2	2	2
Carboxymethyl cellulose	2	2	2	2	2	2
Proximate Composition (g/100g/DM)						
Crude protein	30.0	31.0	31.8	32.0	32.9	33.1
Ether extract	8.0	8.06	8.57	8.70	8.99	8.78
Nitrogen free extract	43.4	42.8	42.0	41.7	41.0	40.1
Crude fibre	1.24	1.22	1.20	1.19	1.17	1.15
Ash	6.64	6.41	6.20	6.08	5.89	5.88
G. Energy (kcal/g/DM)	398.8	398.9	404.0	405.2	409.3	404.7
Protein – Energy Ratio	76.8	77.8	78.6	79.0	80.2	81.7

¹ supplied as (mg/kg diet): Ca-pantothenic acid, 40; pyridoxine, 10; riboflavin, 12; niacin, 20; folic acid, 2; choline chloride, 2000; thiamin HCl 2; D-biotin, 0.28; cyanocobalamin (B₁₂), 0.04; menadione sodium bisulphate (vitamin K), 2; ascorbic acid (vitamin C), 100; DL- α -tocopherol acetate (vitamin E), 100; cholecalciferol (vitamin D), 4000IU; retinyl acetate (vitamin A), 5000IU; salt (NaCl), 6200; copper sulphate (CuSO₄.5H₂O), 56; ferrous sulphate (FeSO₄.7H₂O), 130; manganese sulphate (MnSO₄.H₂O), 178; zinc sulphate (ZnSO₄.H₂O), 120; potassium iodide (KI), 16; corn starch

Table 3: Growth, nutrient utilization and apparent digestibility coefficient ADC) of *O. niloticus* fed tamarind diets for 56 days

Parameters	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Diet 6
Initial mean weight (g)	6.10 ±0.17	6.30 ±0.17	6.36 ±0.18	6.04 ±0.19	6.04 ±0.18	6.00 ±0.17
Final mean weight (g)	13.13±0.52	13.66±0.51	14.23±0.53	14.25±0.51	13.61±0.51	13.04±0.52
Weight gain (g)	7.03 ±0.47	7.30 ±0.46	7.87 ±0.48	8.21 ±0.46	7.57 ±0.46	7.04 ±0.48
SGR (g/day)	1.37 ±0.06	1.37 ±0.05	1.44 ±0.03	1.53 ±0.05	1.45 ±0.06	1.39 ±0.05
FCR	2.29 ±0.09	2.31 ±0.08	2.21 ±0.10	2.08 ±0.07	2.17 ±0.10	2.27 ±0.08
PER	3.67 ±0.24	3.76 ±0.25	3.96 ±0.25	4.10 ±0.24	3.69 ±0.25	3.41 ±0.24
ADC protein (%)	84.9 ±3.05	87.0 ±3.04	89.6 ±3.10	91.2 ±3.11	88.2 ±3.08	82.9 ±3.12
ADC energy (%)	82.4 ±3.41	84.9 ±3.40	87.9 ±3.43	89.7 ±3.42	86.4 ±3.43	80.6 ±4.13

Values along the same row are not significantly different ($P > 0.05$)

Table 4: Carcass composition and blood parameters of *O. niloticus* fed for 56 days

Fish Fed	Crude protein	Ether extract	Ash	TEC (million/il) ^a	PCV (%) ^b	RBC (10 ⁴ mm ³) ^c
Before feeding	36.3	6.25	27.6	3.0	30	180
Diet 1	53.5	11.2	21.5	3.0	29	181
Diet 2	53.6	11.1	21.5	3.1	31	179
Diet 3	53.7	11.2	23.1	3.2	30	182
Diet 4	53.8	11.2	23.9	3.0	31	180
Diet 5	54.0	11.2	23.8	3.1	29	181
Diet 6	54.1	11.6	24.4	3.2	31	180

a. Total erythrocyte count b. Pack cell volume c. Red blood cells

This protein content is higher than that of maize (10%). The major anti-nutrients found in tamarind seed meal were low values of phytin and tannin. Gross composition of the experimental diets, the proximate composition, gross energy (GE) and protein-energy ratio (P: E) were presented in Table 2. The mean value of the dietary fat

recorded from the present study is within the acceptable range recommended for fish culture (Teshima and Kanazawa, 1986). The low fibre contents of the experimental diets depict the high quality which can support high feed digestibility and good fish yield. The mean value of the GE obtained is close to the value of 467 kcal.g⁻¹ used

by Abdelghany (2000) for good tilapia production. Li *et al.* (1991) showed that P:E of diets influenced nutrient utilization and growth; and optimum P:E boosted the profitability of diets and protein sparing ability of the diets (Xiquin *et al.* 1994).

The mean P:E obtained from this study was high but lower than 100mg kcal⁻¹ recorded by Santiago and Laron (1991) as optimum for red tilapia fed 30 % crude protein diets. However, the value compared favourably with the mean value of 86 ± 1.6 which supported good yield of *O. niloticus*, (Nwana and Daramola, 2000). Results of the feeding trials (Table 3) showed no significant variations in the mean weight gain (MWG), specific growth rate (SGR), food conversion ratio (FCR) and protein efficiency ratio (PER) among the fishes fed the control diet and those fed diets supplemented with tamarind seed meal at 20, 40, 60, 80 and 100 % levels of substitution. The mean values of SGR and FCR obtained from this study was close to the mean values of 1.28 ± 0.17 and 1.77 ± 0.33 for SGR and FCR reported for *O. niloticus* fed cottonseed meal based diets Mbahinzireki *et al.* (2001). The SGR obtained was also similar to the mean value of 1.53 ± 0.06 documented for *O. niloticus* fed compounded diets (Maina *et al.* 2002). Furthermore, the mean value of PER obtained from the present study compared well with the mean values of PER of 2.64 ± 0.46 , and 2.48 ± 0.55 reported by Hossain *et al.* (2002) and Ulloa and Verreth (2002) for *O. niloticus* fed dhaincha seed meal based diets and bacteria-treated coffee pulp meal based diets, respectively.

The digestibility of individual ingredients in compounded diet is one of the important factors affecting the growth of fish (De Silva *et al.*, 1996). Uys (1988) noted that rather than look at growth responses, the digestibility of nutrients and energy contents of feedstuffs could be used to assess the suitability and nutritive value of feedstuffs/diets in fishes. The apparent digestibility coefficient (ADC) for protein and energy (Table 3) obtained from the present study were high. There were no significant variations in the ADC for protein and energy of the fishes fed diets 1 – 6 ($p > 0.05$). However, the ADC for protein and energy were marginally higher in fish fed 60 % of tamarind seed meal based diet than in fishes fed other diets ($p > 0.05$). The high values of the ADC for protein and energy recorded from this study indicated that the diets were well digested. The mean values of ADC for protein and energy obtained are close to the ADC for protein (93.0%) and energy (83.0%) recorded by Degani *et al.* (1997) for hybrid tilapia (*O. niloticus*[♂] x *O. aureus*[♀]) fed corn meal. Furthermore, the ADC for protein recorded from the present experiment was similar to mean ADC

value for protein of (81.7 ± 7.3) reported by Mbahinzireki *et al.* (2001).

The carcass composition and blood parameters of the fishes fed diets 1-6 for 56 days are presented in Table 4. Protein and fat deposition were similar in all the fishes fed diets 1–6, and there was no significant difference in ash content ($p > 0.05$). The total erythrocyte count (TEC), pack cell volume (PCV) and the red blood cells (RBC) were high. There were no significant variations in TEC, PCV and RBC of the fishes fed all the diets ($p > 0.05$). The RBC value in this study was close to the range of 700,000-2,000,000 cells mm³ recommended by Saunders (1966) for healthy teleosts. The mean PCV value obtained from the present study was similar to the value of 32.1 % reported by Abdelghany (2000), and the mean value of 36 ± 0.5 reported for *O. niloticus* by Mbahinzireki *et al.* (2001).

ACKNOWLEDGEMENTS

The authors are very grateful to African Academy of Sciences (AAS) Nairobi Kenya for providing the Research Grant No. 99/8 of 3/4/00 used in conducting the study.

REFERENCES

- ABDELGHANY, A. E. (2000). Replacement value of cystine for methionine in semipurified diets supplemented with free amino acids for the Nile tilapia, *Oreochromis niloticus* L. fry. Pages 109 – 119. In: K. Fitzsimmons and J.C. Filho, editors. Proceedings of the Fifth International Symposium on Tilapia Aquaculture. Rio de Janeiro, Brazil, September 3 – 7, 2000.
- AOAC (1990). *Official Methods of Analysis*. Association of Official Analytical Chemist (15th ed.) AOAC Incorporated, Arlington, Virginia, USA. 1094 pp.
- BALOGUN, A. M. (1990). Feeding fish with alternative feed formulations and resources. Pages 669 – 97. In: G. M. Babatunde. (Ed). *Proceedings of the national workshop on alternative feed formulations of livestock and fish feeds in Nigeria*. Economic Affairs Office of the Presidency, Lagos, Nigeria.
- DEGANI, I.G., BIOLA S. and YEHUDA Y. (1997). Apparent digestibility of protein and carbohydrate in feed ingredients for adult tilapia (*Oreochromis niloticus* x *O. aureus*) hybrid. *Bamidgeh*, 49 (3): 15 – 123.
- DE SILVA, S. S., SHIM, K. F and KONG A. K. (1986). An evaluation of the methods used in digestibility estimations of a dietary ingredient and comparison on external, and internal markers, and time

- of faeces collections in digestibility studies in fish *Oreochromis aureus* (Steindacher). *Reproduction, Nutrition and Development*, 30: 215 – 226.
- FAO (1970). *Amino acid content of foods and biological data on protein*. FAO Nutritional studies, Number 24, Food and Agriculture Organisation, Rome, 46 pp
- HALVER, S. E., YIMAN, A. and SMITH, R. R. (1993). Acid insoluble ash as a convenient method for estimating digestibility component in Diets. Abstract, *International Conference for World Agriculture*, Fresh Water Fish Poster, pp 230.
- HARRIS, L. E. (1970). *Nutrition Research Techniques for Domestic and Wild Animals*. Vol. 1, pp 84321. Utah
- HOSSAIN, M. A., FOCKEN, U. and BECKER K. (2002). Nutritional evaluation of dhaincha (*Sesbania aculeata*) seeds as dietary protein source for tilapia *Oreochromis niloticus*. *Aquaculture Research*, 33: 653 – 662.
- LI, Z., YE, W. and HE, X. (1991). The nutritional value of commercial feed ingredients for Nile tilapia (*Oreochromis niloticus*) in China. Pages 101 – 106. In: S.S. De Silva, Editor. *Fish nutrition research in Asia*. Proceedings of the fourth Asian Fish Nutrition Workshop. Asian Fish Society Special Publication, Asian Fisheries Society, Manila, Philippines.
- MBAHINZIREKI, G. B., DABROWSKI, K., LEE, K. J., EL-SAIDY, D. and WISNER, E. R. (2001). Growth, feed utilization and body composition of tilapia (*Oreochromis* sp.) fed with cottonseed meal based diets in a recirculating system. *Aquaculture Nutrition*, 7: 189 – 200.
- MAINA, J.G., BEAMES, R. M., HIGGS, D., MBUGUA, P. N., IWAMA, G. and KISIA S. M. (2002). Digestibility and feeding value of some feed ingredients fed to tilapia *Oreochromis niloticus* (L). *Aquaculture Research*, 33: 853 – 862.
- NWANNA, L. C. and DARAMOLA, J. A. (2000). Harnessing of shrimp head waste in Nigeria for low cost production of Nile tilapia (*Oreochromis niloticus* L). Pages 174 – 178. In: K. Fitzsimmons and J.C. Filho, Editors. *Proceedings of the Fifth International Symposium on Tilapia Aquaculture*. Rio de Janeiro, Brazil, September 3 – 7, 2000.
- OLIVERA-NOVOA, M.E., CAMPOS, G. S., SABIDO G. M. and MARITINEZ-PALACIOS, C. A. (1990). The use of alfalfa leaf protein concentrates as a protein source in diets for tilapia *Oreochromis niloticus*. *Aquaculture*, 9: 91 – 302.
- SANTIAGO, C. B. and LARON M. A. (1991). Growth and carcass composition of tilapia fry fed diets with varying protein levels and protein to energy ratios. Pages 55 – 62. In: S. S. De Silva, Editor. *Fish nutrition research in Asia*. Proceedings of the fourth Asian Fish Nutrition Workshop. *Asian Fish Society Special Publication*, Asian Fisheries Society, Manila, Philippines.
- SATHE, S. K. and SALUNKHE, D. K. (1981). Investigations on winged beans (*Psophocarpus tetragonolobus* (L) proteins and antinutritional factors. *Journal of Food Science*, 46: 1389 – 1392.
- SAUNDERS, A. C. (1966). Elasmobranch blood cells. *Copeia*, 196: 348 – 351.
- SVOBODOVA, Z., PRAVDA, D. and PALACKOVA, J. (1991). *Unified methods of haematological examinations of Fish*. Research Institute of Fish Culture and Hydrobiology, Vodany, Czechoslovakia, 49 pp.
- TACON, A. G. J. (1993). *Feed ingredients for warm water fish: fish meal and other processed feedstuffs*. FAO Fisheries Circular, No. 856, FAO, Rome, 64 pp.
- TESHIMA, S. and KANAZAWA, A. (1986). The growth and nutritive requirements of tilapia *Journal of Aquaculture Japan*, 23: 106 – 111.
- ULLOA, J. B. and VERRETH, J. A. (2002). Growth, feed utilization and nutrient digestibility in tilapia fingerlings (*Oreochromis aureus* Steindachner) fed diets containing bacteria-treated 233. coffee pulp. *Aquaculture Research*, 33: 189 – 195.
- UYS, W (1988). Breeding of African catfish, *Clarias gariepinus*. Pages 47 – 61. In: Hecht, T. W. Uys and P.J. Britz (eds.). *The culture of sharptooth catfish Clarias gariepinus in Southern Africa*. South African National Scientific Programme Report No.153, South African.
- XIQUIN, H., LIZHU, J., YUNXIA, Y. and GUOHAUN X. (1994). Studies on the utilization of carbohydrate – rich ingredients and optimal protein energy ratio in Chinese Bream, *Megalobrama amblycephala* Yin. Pages 31 – 42. In: S. S De Silva, Editor. *Fish Nutrition Research in Asia*. Proceedings of the Fifth Asian Fish Nutrition Workshop. Asian Fisheries Society Special Publication, Number 9, Manila, Philippines, Asian Fisheries Society.
- ZAR, J. H. (1984). *Biostatistical analysis*. Prentice Hall, Englewood Cliffs, NJ USA. 383 pp.

