

***A  
N  
I  
M  
A  
L***



***R  
E  
S  
E  
A  
R  
C  
H***



Volume 4 Number 2 (2007)

***I  
N  
T  
E  
R  
N  
A  
T  
I  
O  
N  
A  
L***



**An International Journal Publishing Original Research Involving  
the Use of Animals and Animal Products**

**ISSN: 159-3115**

**Website: [zoo-umn.org](http://zoo-umn.org)**

## MOLLUSCAN POPULATION OF AN AFRICAN ARID ZONE LAKE

IDOWU Rachel Toyosi, GADZAMA Usman Ngamarju, ABBATOIR Ahmed and INYANG Nicholas Mathias

<sup>1</sup>Department of Biological Sciences University of Maiduguri, Maiduguri, Borno State, Nigeria

<sup>2</sup>Department of Zoology, Faculty of Biological Sciences University Nigeria Nsukka

**Corresponding Author:** Idowu, R. T. Department of Biological Sciences University of Maiduguri, Maiduguri, Borno State, Nigeria. Email: [toyosi\\_idowu@yahoo.com](mailto:toyosi_idowu@yahoo.com) Phone: 2348023579099

### ABSTRACT

*The species composition, abundance and distribution of molluscs population together with some physico-chemical variables from five different stations in the littoral region of lake Alau, Maiduguri; Borno state, were studied from October 2001 to September 2002. Three patterns of seasonal abundance were found, maximal abundance during the rainy season (July -September), moderate abundance during the harmattan season (November – February) and minimal abundance in the dry hot season (March - June). Significant differences in species composition and abundance of the molluscs were found between stations studied. The total number of organisms recorded was 3368 comprising of 1924 Bivalves and 1544 Gastropoda. 8 families were recorded which includes Bithyniidae, Hydrobiidae, Lymnaeidae, Physidae, Valvatidae, Vivipariidae, Sphaeridae and Unionidae while 15 species were observed in this study. The predominant families in terms of total number collected from all stations Sphaeridae and Unionidae recording 1006 and 933 organisms with the percentage compositions of 29.86% and 27.7%. The least was from the families Lymnaeidae with total number of 116 and 3.44% as the percentage composition. The abundance of the molluscs was positively and significantly correlated at 5% confidence with temperature ( $r = 0.675$ ), dissolved oxygen ( $r = 0.832$ ), phosphate ( $r = 0.528$ ).*

**Keywords:** Molluscs, Littoral Region, Abundance, Unionidae

### INTRODUCTION

The Phylum Mollusca inhabits permanent water bodies across a large range in Africa. Previous work includes studies by Appleton (1974, 1977) on the Mollusca composition of pond and river in South Africa, Babiker *et al.* (1985) on the snails of irrigation system in Sudan, Woolhouse and Chandiwana (1989) on River Zimbabwe snail composition, Akuforgwe *et al.* (1995) Mollusca of Jos dam, Imafidon, (1991); and Omudu and Iyough (2005) on River Benue Mollusca. These studies suggest that the population dynamics of the phylum was greatly affected by the water chemistry.

Idowu (2004) observed that shallow Lake often with a well developed littoral vegetation are often more productive than deep lakes and particularly in the tropics, they are important source for fisheries and other aquatic fauna products. The biology of molluscs has a status which allows a great deal of environmental exploitation because of its relationship with its environment such as being commercial important as source of income, nutrients for certain organism, as well its ability to exist as host of certain parasites of man (Dussart, 1977). These and certain other relationship have greatly necessitated the need for study of the organism. In this paper ecological studies of population dynamics of molluscs in Lake Alau are reported. The data include the study of some physical chemical characteristic as well as relationship between molluscs abundance and these variables.

### MATERIALS AND METHODS

Lake Alau is one of the several tropical lakes in Africa, and it is the second largest lake in Borno state, Nigeria. Lake Alau was created in 1987 by damming river Ngadda about 22 km from Maiduguri, along Bama road. It is located between latitude 13° N and 14° N, and longitude 12° E and 13° E. It has a total surface area of 56 km<sup>2</sup> (CBDA, 1986, Bankole *et al.*, 1994). Being located in the north – east arid zone, the climate is Sahelian with three distinct seasons. The rainy season starts from June to October, the harmattan season with dry cold wind from November to February and dry hot season with extreme temperature from March to May. It has a mean depth of 9.5 m, temperance of about 0.48 m, total alkalinity of 38.4 mg/l (Idowu, 2004; Idowu *et al.*, 2004). The study was carried out for over a 12 months period from October 2001 to September 2002. Five stations were chosen for this study based on accessibility, fishing activities, irrigation and drinking spot for animals. The description of each station is as described in Idowu *et al.*, (2004).

The stations were sampled fortnightly using 0.2 x 0.2 m scoop with a mesh size 2 x 2 mm. Fabricated serrated edge cylindrical bucket of diameter 50 cm and 30 cm height were also used in collection of samples. Each station was sampled in blocks of 10 m lengths. The samples were sorted in the Department of Biological Sciences Laboratory, University of Maiduguri into macro and micro samples using a sieve of 9 meshes per cm.

Each specimen was identified to species level using the keys of Mocas (1959), Brown (1970), Pennak (1978), Fitter and Manuel (1986). The relative abundance and dominance of each group was obtained by direct ratio comparisons using Sorenson diversity index (Margalef, 1982); Shannon (1948) Index as modified by Wilhm (1975) was used to characterize the integral richness and the eaves of distribution.

Qualitative and quantitative data on selected water quality were measured during each visit to the sites. These include water temperature, depth, dissolved oxygen, pH, water current, biochemical oxygen demand (BOD) and conductivity (Boyd, 1979; Apha, 1989). All data on the physical, chemical and biological studies were assessed for normality and homogeneity of variance. Correlation coefficient in relation to gastropod distribution was determined. Data collected for gastropod were subjected to two-way analysis of variance (ANOVA) and F - LSD

## RESULTS

The mollusc population in Lake Alau comprised of 2 classes (Gastropoda and Bivalva), 8 families and 15 species (Table 1). Gastropoda had 6 families and 9 species, while Bivalva had 2 families and 6 species respectively. Sphaeriidae and Unioniidae had the highest species composition. Hydrobiidae and Valvatidae had 2 species each. All other families i.e. Bithyniidae, Physidae, Vivparidae has a species each.

Most of these species were distributed in all the station except *Lampris radiates*, *viviparous species*, and *Bithynia tentaculata*, that were not found in stations 1 and 2. *Valvata sincera* was also absent in station 2 and 3. All other families were represented in all the stations, except Bithyniidae, Vivpariidae and Unionidae (Table 1).

The percentage composition calculated for each family shows that Sphaeriidae had 29.86%, followed by Unionidae (27.7 %), Bithyniidae (10.90 %) and Physidae (9.85%) respectively. There were no significant difference ( $P > 0.05$ ) between Sphaeriidae and Unionidae. No significant difference ( $P < 0.05$ ) was also observed between Bithyniidae and Physidae.

The distribution of molluscan population in relation to stations (Table 2) showed that station 4 had the highest number of species composition (15 species) and all the species were present in this station. The percentage composition in relation to stations showed that station 4 had 48.9% followed by station 5 with (17.5 %), with 14 species. Station 2 had 8.7% with 11 species (Table 1 and 2).

The Sorenson's index of similarity between stations showed the degree of similarity in this order, station 4 (89.96) higher than station 5 (76.66) and stations 3, 1 and 2 with 69.28, 58.50 and 52.72 similarities respectively. No significant difference ( $P > 0.05$ ) was observed between molluscan similarities of stations.

The monthly variation in the population and abundance of Mollusca classes showed that Gastropoda were the most abundant group, and were dominant in all the months except between July and September.

Highest population abundance was observed between July and September (rainy season) which was significantly different ( $P < 0.05$ ) from all other months. A drastic decline in total abundance of both classes was observed between February and May (dry hot season). However, there was an increase in the total numbers collected from October to January (Harmattan season). The seasonal variations showed 3 periods i.e. population increase (October to January). Maximal abundance (June to September) and population decrease (February to May). Gastropoda and Bivalva were found to exhibit the same seasonal periodicity, with marked differences in the relative abundance of species in the various months.

The physicochemical parameters of the five sampling stations are summarized on Table 3. Water temperature varied between  $25.05 \pm 0.4^\circ \text{C}$  and  $27.24 \pm 0.12^\circ \text{C}$ , the current was between  $19.62 \pm 0.30 \text{ cm/sec}$ . The highest mean value for transparency was  $0.42 \pm 0.03 \text{ m}$ , while  $0.26 \pm 0.01 \text{ m}$  was the lowest. The dissolved oxygen varied between  $5.15 \pm 0.03 \text{ mg/l}$  and  $6.35 \pm 0.05 \text{ mg/l}$ .

The result of the correlation coefficient calculated between selected physicochemical characteristics and molluscs abundance showed a significant positive correlation with water temperature (0.675), dissolved oxygen (0.832). pH (0.710), current 0.528 and conductivity 0.899 (Table 4).

## DISCUSSION

The seasonal variations of molluscs population may provide hints as to the extent of environment perturbation as the populations were proportionally higher in the rainy season and harmattan season. The same strong and pronounced seasonality of macroinvertebrates was observed by Dejoux *et al.* (1971) and Mbagwu (1993) in lake Chad and Tiga respectively. Also, discernable seasonal changes have been recorded in sub tropical Lake Sibaya (Hart, 1993), and Okomu forest reserve sanctuary in Nigeria Ogbaisu *et al.* (1995).

Climatic regime has long been known to explain variation in distribution pattern of aquatic invertebrates including molluscs, and thus community structure at global taxa. It was observed that mollusc fauna were higher in the rainy season especially for Gastropoda. It is likely that the rainy periods which give organisms more opportunity to colonize different habitats contribute to the abundance. The cooler and wetter environment may simply be a more suitable environment for the species

The abundance of gastropod in lake Alau may be due to the types of aquatic habitats in and around the Lake. It is also possible that Gastropoda abundance is a consequence of difference in protection and avoidance from the predators. It is unlikely that fish affect Gastropod distribution in lake Alau.

The overall composition and abundance of mollusc family and species in this study varied both spatially and temporally in response to selected physical chemical factors of the aquatic environment.

**Table 1: Molluscan distribution in relation to stations in Lake Alau**

Phylum	Class	Family	Species	Station				
				1	2	3	4	5
Mollusca	Gastropoda	Bithyniidae	<i>Bithynia tentaculata</i>	-	-	+	+	-
		Hydrobiidae	<i>Potamopyrgus jenkinsi</i>	+	+	-	+	+
			<i>Hydrobiidae immatures</i>	+	+	+	+	+
		Lymnaciidae	<i>Lymnae truncatula</i>	+	+	+	+	+
			<i>Lymnae palustris</i>	+	+	+	+	+
		Physidae	<i>Physelia species</i>	+	+	+	+	+
		Valvatidae	<i>Valvata lewisi</i>	+	+	+	+	+
			<i>Valvata sincera</i>	+	-	-	+	+
		Viviparidae	<i>Viviparous species</i>	-	-	+	+	+
		Bivala	Hydrobiidae	<i>Pisidium casternum</i>	+	+	+	+
	Physidae		<i>Pisidium nitidum</i>	+	+	+	+	+
	Sphaeridae		<i>Sphaerum nitidum</i>	+	+	+	+	+
	Unionidae		<i>Unio species</i>	+	+	+	+	+
	<i>Elliptia campalanata</i>		+	+	+	+	+	
			<i>Lampris radiate</i>	-	-	-	+	+
<b>Total</b>	<b>2</b>	<b>8</b>	<b>15</b>	<b>12</b>	<b>11</b>	<b>12</b>	<b>15</b>	<b>14</b>

Key: - Absent, + Present

**Table 2: The total abundance and percentage composition of Mollusca families collected in relation to station in Lake Alau**

Taxa	Stations					Total Collected	Percentage composition
	1	2	3	4	5		
Bithyniidae	-	-	106	260	-	366	10.90
Hydrobiidae	63	50	25	100	45	283	8.40
Lymnaciidae	10	13	10	48	35	116	3.44
Physidae	46	30	22	200	34	332	3.85
Valvatidae	10	15	18	55	35	133	3.95
Viviparidae	-	-	46	88	65	199	5.9
Sphaeridae	136	168	150	390	162	1006	29.86
Unionidae	39	20	162	500	212	399	27.70
<b>Total</b>	<b>304</b>	<b>296</b>	<b>539</b>	<b>1641</b>	<b>588</b>	<b>3368</b>	<b>100</b>
<b>%/station</b>	<b>9.02</b>	<b>8.78</b>	<b>16.0</b>	<b>48.7</b>	<b>17.5</b>		<b>100</b>

**Table 3: Physico-chemical parameters in relation to stations in lake Alau**

Parameters	1	2	3	4	5
Temperature (°C)	25.25 ± 0.18 <sup>b</sup>	25.05 ± 0.14 <sup>b</sup>	27.24 ± 0.19 <sup>b</sup>	27.24 ± 0.12 <sup>a</sup>	25.13 ± 0.00 <sup>b</sup>
Current (cm/sec)	26.71 ± 0.30 <sup>b</sup>	25.46 ± 0.27 <sup>b</sup>	25.08 ± 0.36 <sup>b</sup>	25.10 ± 0.28 <sup>b</sup>	19.62 ± 0.22 <sup>a</sup>
Transparency (m)	0.36 ± 0.01 <sup>b</sup>	0.33 ± 0.02 <sup>b</sup>	0.35 ± 0.01 <sup>b</sup>	0.42 ± 0.03 <sup>a</sup>	0.26 ± 0.01 <sup>c</sup>
PH	6.79 ± 0.05 <sup>b</sup>	6.97 ± 0.02 <sup>b</sup>	6.83 ± 0.02 <sup>b</sup>	7.29 ± 0.05 <sup>b</sup>	6.59 ± 0.01 <sup>b</sup>
Dissolved oxygen (mg/l)	6.15 ± 0.05 <sup>a</sup>	6.35 ± 0.05 <sup>a</sup>	5.18 ± 0.02 <sup>b</sup>	6.32 ± 0.01 <sup>a</sup>	5.15 ± 0.03 <sup>b</sup>
Biochemical Oxygen Demand(mg/l)	4.34 ± 0.32 <sup>a</sup>	4.30 ± 0.28 <sup>a</sup>	4.45 ± 0.50 <sup>q</sup>	5.03 ± 0.33 <sup>a</sup>	5.31 ± 0.25 <sup>a</sup>
Conductivity (ohms/cm)	131.45 ± 0.75 <sup>b</sup>	128.45 ± 0.52 <sup>b</sup>	119.42 ± 0.83 <sup>a</sup>	115.47 ± 0.75 <sup>a</sup>	118.47 ± 0.16 <sup>a</sup>

**Table 4: Correlation Coefficient for selected physical and chemical parameters and mollusc abundance in lake Alau**

Physical/ chemical Parameters	Correlation Coefficient "r" P < 0.05
Water temperature	0.675
Dissolved oxygen	0.832
Phosphate	0.648
Ph	0.710
Current	0.528
Conductivity	0.899
BOD	

The overriding influence of the temperature, current, dissolved oxygen, conductivity, biochemical oxygen demand (BOD) in distribution and abundance can explain the significant lower numbers observed in all stations between February and June. The effect of the rainy season between July and September may have increased feeding habitat and access to breeding. The various ecological requirements and

also water quality parameters affected the spatial distribution and abundance. Various physical chemical factors collectively have an effect on the abundance of molluscs under condition. Okafor (1990) explained how rainfalls affect the quality of the habitat making it suitable or unsuitable for the molluscan population and abundance.

The seasonal pattern of the total mollusc abundance is consistent with the observation of Obureke (1980), Okafor (1990), Omudu and Iyough (2005). This seasonal dynamic is attributed to seasonal periodicity of the quality and quantity of edible, competition, resumption of normal metabolic activities, by those that have gone through period of adverse conditions, interaction as well as climatic changes in the natural environment. The distribution and abundance of the molluscan population in lake Alau may also be attributed to the availability of food, shelter and oviposition sites. This agrees with Whitton 1975, Omudu and Iyough (2005) that water bodies rich in organic and silt matter are known to support thriving populations of macroinvertebrates. The dominant species encountered in this study i.e. *Hydrobiidae* immature, *Lymnaea palustris*, *Physelia* species.; *Valvata lawisi*, *Pisidium casertanum*, *Pisidium nitidum*, *Sphaerium nitidum*, *unio* species and *Eliptia campalanata* were encountered in all the station surveyed.

The correlation coefficient value between the selected physical and chemical parameters and the molluscs abundance showed significant correlation. This agrees with the findings of Imafidon (1991), Agi (1995), Okafor and Ngang (2004) that fresh water molluscan populations thrive well on environment with good high water qualities.

In conclusion, the temperature appears to be the most important factor influencing the development and distribution of molluscs in the lake Alau. Its effect may be direct in presenting optimum conditions for chemical activities for molluscan population, and indirect in distribution within the habitats. Arad *et al.* (1992, 1993), Okafor (1991), Idowu *et al.* (2005) suggested that climate and microhabitats are the main determinants of species resistance to desiccation, availability, distribution and abundance of molluscs in fresh water environment. The present study suggests that these factors may likewise play important roles in their distribution.

## REFERENCES

- AGI, P. I. (1995). Survey of Freshwater Snail Vectors of Schistosomiasis and study of Physico chemical parameters of the water bodies in Ogoni Communities, River State, Nigeria. *Acta Hydrobiologica*, 21: 3 – 36
- AKUFORGWE, P. F., DONDJJI, B., OKWUOSA, U.N., DAKUL, D. A and NTONIFOR, H. N. (1995). Observed disparity on Schistosome infection rates in field *Biomphalaria pfeifferi* between two areas of the Jos metropolis, Nigeria. *Parasites*, 2: 89 – 91.
- APHA, A. (1979). *Standard Methods for the examination of water and waste water*, 17<sup>th</sup> Edition. American Public Health Association, Washington D.C. N.Y. 1134 pp.
- APPLETON, C. C. (1974). The population fluctuation of five freshwater snails in the eastern Transverse lowland and their relationship to known bilharzias transmission patterns, *South African Journal of Science*, 70: 114 – 150.
- APPLETON, C. C. (1977). The influence of temperature on the life cycle and distribution of *Biomphalaria pfeifferi* (krauss, 1948) in South-Eastern Africa. *International Journal of Parasitology*, 7: 335 – 345.
- ARAD, Z., GOLDENBERG, S., and HELLER, J. (1992). Intraspecific variation in resistance to desiccation and climate gradients in the distribution of the land snails *Xeropicta vestalis*. *Journal of Zoology*, 226: 643 – 656
- ARAD, Z., GOLDENBERG, S. and HELLER, J. (1993). Intraspecific variation in resistance to desiccation and climate gradients in the distribution of the land snails Trochoidae Simulata. *Journal of Zoology*, 229: 249 –265
- BABIKER, A., FENWICK A, DAFFALLA, A. A. and AMIN, M. A. (1985). Focality and Seasonality of *Schistosoma mansoni* transmission in the Gezira Irrigated Area, Sudan. *Journal of Tropical Medicine and Hygiene*, 88: 57 – 63
- BANKOLE, N. O., SULE, O. D., OKWUUNDU, E. C. and AMADI, M. (1994). Preliminary Investigation into the fish and Catch assessment survey of Alau Lake. *Annual Report of National Institute for Freshwater fisheries research (NIFFR)* 28 pp.
- BOYD, C. E. (1979). *Water quality in warm water fish pond*. Craft Master prints Incorporated, Auburn, Alabama. 353 pp.
- BROWN, A. L. (1970). *Key to pond organisms*. Nuffield Advanced Science Penguin Books Limited, England.
- CBDA (1986). A resettlement plan for the Lake Alau dam and Jere bowl scheme. Agricultural survey and background studies. *Report submitted to Chad Basin Development Authority by Askon Nigeria Limited*. 59 pp
- DEJOUX, C., LAUZANE, L. and LEVEQUE, C. (1971). Nature des fords et repartitiondes organismes benthiques don la region de bal (Lac T. Chad) Cahiers ORSTOM series. *Hydrobiologia*, 5: 213 – 223
- DUSSART, G. B. J. (1977). The ecology of *Potamopygus jenkinsi* (smith) in North West England with a note on Marstoniopsis Schottzi (Schmidt). *Journal of Molluscan Studies*, 43: 208 – 216
- FITTER, R. and MANUEL R. (1986). *Field guide to fresh water life*. William Collins and Son Toronto. 382 pp.
- HART, H. I. (1993). A national approach to river. *Search*, 24: 125 –130
- IDOWU, R. T. (2004). *Limnological Studies of Lake Alau, Borno State, Nigeria*, PhD Thesis, Department of Zoology, University of Nigeria Nsukka, Nigeria 189 pp.
- IDOWU, R. T., INYANG N. M, and EYO, J. E. (2004). The physical chemical parameters of an African Arid Zone man made Lake. *Animal Research International*. 1(2): 113 – 119

- IDOWU, R. T., INYANG N. M., and MGBENKA, B. O. (2005). Macroinvertebrate studies of the littoral region in an arid Zone Lake Alau, Maiduguri, Borno State. *Pecop Resources*, 11: 13 –18.
- IMAFIDON, E. U. (1991). Ecological studies of Fresh water snails in Ibadan, Nigeria, *Nigerian Journal of Parasitology*, 12: 59 – 63.
- IMAFIDON, E. U. and IYOUGH, A. (2005). Ecological Studies of fresh water snails in Ibadan, Nigeria. *Nigerian Journal of Parasitology*, 12: 59 – 63.
- MACAN, T. T. (1959). *A guide to freshwater Invertebrate animals*. Longman England. 406 pp.
- MARGALEF, R. (1982) Phytoplankton composition and distribution a san expression of properties of reservoirs. *Canadian Water Resources Journal*, 7: 26 50.
- MBAGWU, I. G. (1993). Vertical distribution of macrobenthos invertebrates in the profundal sediments of the impoundment of the dammed river Dambe; *Australian Journal of Aquatic Science*, 5: 19 – 25.
- OBUREKE, G. T. (1980). *Studies on the feeding biology anatomical variation and ecology of vector of Schistosomiasis and other fresh water Snails in South western Nigeria*. PhD Thesis, University of Ibadan 370 pp.
- OGBEIBU, A. E., EGBORGE, E. and VICTOR, R. (1995). Hydrobiological studies of water bodies in the Okomu forest reserve (Sanctuary) in the Southern Nigeria; distribution and diversity of the invertebrates fauna. *Tropical Freshwater Biology*, 4: 1 – 27.
- OKAFOR, F. C. (1990). *Schistosoma haematobium* cercariae transmission pattern in Freshwater systems of Anambra State Nigeria, *Angewandte Parasitologie*, 31: 159 –166.
- OKAFOR, F. C. (1991). The effect of temperature on embryonic development and reproduction of *Bulinus* (Physopsis) *globosus* (Morelet) and *Lymnaea natalensis* (Krauss) (Gastropoda, Pulmonate). *Journal of Aquatic Science*, 6: 7-11
- OKAFOR, F. C. and NGANG, I. (2004). Freshwater Snails of Niger-cem, Nkalagu Eastern Nigeria, Observations on some Demographic aspects of the Schistosome-transmitting Bulinids. *Animal Research International*, 2: 120 – 124.
- OMUDU, E. A. and IYOUGH A. (2005). Ecological Studies of the Gastropod fauna of some minor tributaries of River fauna. Makurdi, *Nigeria Animal Research International*, 2: 306 – 310.
- PENNAK, R. W. (1978). *Freshwater Invertebrate of the United States* 2<sup>nd</sup> Ed. Wiley Inter, Science, New York 265 pp.
- SHANNON, C. E. (1948). A mathematical theory of communication. *Bell Systematic technical journal* 27: 379 – 423.
- WOOLHOUSE, M. E. J. and CHANDIWANA, S. K. (1989). Spatial and temporal heterogeneity in the population dynamic of *Bulinus globosus* and *Bomphalaria pfeifferi* and the epidemiology of their infection with schistosomes, *Parasitology*, 98: 21- 34.
- WHITTON, B. A. (1975). Zooplankton and Macro invertebrates. Pages 87 –118 In: WHITTON, B.A. (Ed) *Studies in River Ecology* vol. 2. Publisher Limited London.
- WILHM, J. L. (1975). Biological indicators of Pollution. Pages 375 – 403. In: WHITTON, B. A. (Ed) *Studies in River ecology* Volume 1. Blackwell Scientific Publication, Oxford.